

Variety Name: **Aleksandra**

Crop: Barley

Breeders Reference: BOLB17.4211

AFP No:

Breeder: Agrovista UK Ltd

Translated from: Polish

Origin:

Date: 06 April 2023

Code: ALEK

*Plant - growth habit:	Intermediate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Early to medium
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Weak
*Ear- glaucosity:	Weak
Ear- attitude:	Semi-erect
Ear- length:	Short to medium
*Ear- number of rows:	Two
*Ear- density:	Lax
Ear- shape:	Tapering to parallel
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Medium
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	N/A
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn cf grain:	Slightly longer
Lemma nerves- anthocyanin colouration:	
*Plant height:	
Collar type:	
Rachis- length of first segment:	Long
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Absent or very weak
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

19 April 2023

Variety Name: **AMISTAR**

Crop: Winter barley

Breeders Reference: MH 05 AM 04

AFP No:

Breeder: SARL

Translated from: French

Origin:

Date: 27 March 2018

Code: AMIS

*Plant - growth habit:	Intermediate to semi prostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Early
*Flag leaf- anthocyanin colouration of auricles:	Medium to strong
Frequency of plants with recurved flag leaves:	Medium
Flag leaf- glaucosity of sheath:	Strong
*Ear- glaucosity:	Medium
Ear- attitude:	Erect to semi erect
Ear- length:	Medium to long
*Ear- number of rows:	more than two
*Ear- density:	Medium to dense
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Medium
*Awn- length compared to ear:	Medium to long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn cf grain:	Equal
Lemma nerves- anthocyanin colouration:	
*Plant height:	Medium
Collar type:	
Rachis- length of first segment:	Short to medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Absent or very weak to weak
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Strong
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Medium to strong
*Grain- ventral furrow - presence of hairs:	Present
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

28 March 2018

Winter Barley

[AFP 2/3597]

AMPLE

CHARACTERS IN DETAIL

Grain

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	medium
Husk (presence after harvest)	husked
Ventral furrow hairiness	absent
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	horizontal
Time of ear emergence	medium to late
Flag leaf glaucosity	strong
Ear glaucosity	medium
Ear attitude (21 days after ear emergence)	horizontal to semi-drooping
Length (stem, ears and awns)	short to medium

Anthocyanin colouration

Flag leaf auricles	absent or very weak
Awn tips	absent or very weak
Lemma nerves (at early dough)	absent or very weak

Ear

Number of rows	two
Shape	slightly tapering
Density	sparse to medium
Length (excluding awns)	long
Awn length (compared to ear)	medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	not applicable

Rachis

Length of first segment	long
Curvature of first segment	medium to strong



ARETHA**CHARACTERS IN DETAIL****Grain**

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	strong
Husk (presence after harvest)	husked
Ventral furrow hairiness	absent
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	horizontal to semi-reflexed
Time of ear emergence	medium
Flag leaf glaucosity	medium to strong
Ear glaucosity	medium
Ear attitude (21 days after ear emergence)	semi-erect to horizontal
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	strong
Awn tips	medium
Lemma nerves (at early dough)	strong

Ear

Number of rows	two
Shape	slightly tapering
Density	sparse to medium
Length (excluding awns)	medium
Awn length (compared to ear)	short to medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	n/a

Rachis

Length of first segment	short to medium
Curvature of first segment	weak

BANSHEE**CHARACTERS IN DETAIL****Grain**

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	absent
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	absent
Flag leaf attitude	semi-erect to horizontal
Time of ear emergence	medium to late
Flag leaf glaucosity	medium
Ear glaucosity	weak
Ear attitude (21 days after ear emergence)	horizontal
Length (stem, ears and awns)	medium to long

Anthocyanin colouration

Flag leaf auricles	strong
Awn tips	strong
Lemma nerves (at early dough)	strong to very strong

Ear

Number of rows	two
Shape	slightly tapering
Density	medium
Length (excluding awns)	long
Awn length (compared to ear)	short to medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	not applicable

Rachis

Length of first segment	long
Curvature of first segment	medium

BOLIVIA**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	long
Shape	tapering

Awn

Length compared to ear	short to medium
------------------------	-----------------

Rachis

Length of first segment	medium
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	husked
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	strong
Awn tips	medium to strong

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	absent
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak
Ear - attitude	semi-erect

BOLTON**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax to medium
Length	medium
Shape	tapering

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	medium
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	very short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak
Ear - attitude	semi-erect

Variety Name: **Bordeaux**

Crop: Winter Barley

Breeders Reference: NOS 911.016-53

AFP No:

Breeder: Nordis Seed A/S

Translated from: Danish

Origin:

Date: 04 May 2020

Code: BORD

*Plant - growth habit:	Intermediate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Medium
*Flag leaf- anthocyanin colouration of auricles:	Very strong
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Strong
*Ear- glaucosity:	Medium to strong
Ear- attitude:	Semi erect to horizontal
Ear- length:	Medium
*Ear- number of rows:	T
*Ear- density:	Two
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Strong
*Awn- length compared to ear:	Short to medium
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	NA
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn cf grain:	Equal
Lemma nerves- anthocyanin colouration:	Strong
*Plant height:	Medium
Collar type:	
Rachis- length of first segment:	Medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Strong
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Absent or very weak
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Strong
*Grain- ventral furrow - presence of hairs:	Present
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

06 May 2020

BUCCANEER

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	husked
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	medium
Lemma nerves - at early dough	weak
Awn tips	weak to medium

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	weak to medium
Ear - attitude	erect

CALIFORNIA

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	long
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	absent
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

CARAT**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax to medium
Length	medium
Shape	tapering

Awn

Length compared to ear	long (longer than ear)
------------------------	------------------------

Rachis

Length of first segment	short to medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	
Lemma nerves - at early dough	absent or very weak
Awn tips	

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong to very strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	horizontal to semi-recurved

CARROUSEL

CHARACTERS IN DETAIL

Grain

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	strong
Husk (presence after harvest)	husked
Ventral furrow hairiness	absent
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	semi-reflexed
Time of ear emergence	medium
Flag leaf glaucosity	medium to strong
Ear glaucosity	medium
Ear attitude (21 days after ear emergence)	horizontal
Length (stem, ears and awns)	medium to long

Anthocyanin colouration

Flag leaf auricles	medium to strong
Awn tips	strong to very strong
Lemma nerves (at early dough)	strong

Ear

Number of rows	six
Shape	parallel
Density	medium
Length (excluding awns)	short
Awn length (compared to ear)	long
Median spikelet - length of glume+awn cf grain	shorter

Sterile Spikelet

Development	not applicable
Attitude	not applicable

Rachis

Length of first segment	medium
Curvature of first segment	very weak

CASSATA**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium to dense
Length	medium
Shape	tapering

Awn

Length compared to ear	+/- equal to longer
------------------------	---------------------

Rachis

Length of first segment	short to medium
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	parallel
Sterile spikelet - development	full
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	absent (grains thresh free)
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	weak
Awn tips	medium to strong

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

CRAFT**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	weak to medium
Lemma nerves - at early dough	strong
Awn tips	strong

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	semi-erect

ELECTRUM**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	weak
Lemma nerves - at early dough	weak to medium
Awn tips	weak to medium

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	early to medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude	erect to semi-erect

FLAGON**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax to medium
Length	long
Shape	parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	horizontal to semi-recurved

FUNKY**CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	dense
Length	short
Shape	parallel

Awn

Length compared to ear	long
------------------------	------

Rachis

Length of first segment	short
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn cf grain	longer

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	strong
Ear - attitude 21 days after ear emerg.	semi-erect

Variety Name: **Idilic**

Crop: **Alternative**

Breeders Reference: sc 50733 rh

AFP No:

Breeder: Secobra

Translated from:

Origin:

Date: 11 May 2022

Code: IDIL

*Plant - growth habit:	Intermediate to semi-prostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Early to medium
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	Horizontal to semi-drooping
Flag leaf- glaucosity of sheath:	Strong
*Ear- glaucosity:	Medium to strong
Ear- attitude:	Horizontal
Ear- length:	Long
*Ear- number of rows:	Two
*Ear- density:	Medium to dense
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Medium
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	Non or rudimentary
Sterile spikelet-shape of tip:	NA
Sterile spikelet- length of lemma:	NA
Median spikelet- length of glume + awn cf grain:	Equal
Lemma nerves- anthocyanin colouration:	Absent or very weak
*Plant height:	Short to medium
Collar type:	
Rachis- length of first segment:	Medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium to strong
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Strongly coloured
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Weak to medium
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Present
*Grain- husk:	PresentClasping
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	
*Seasonal type:	Alternative
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

11 May 2022

INTEGRAL**CHARACTERS IN DETAIL****Grain**

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	strong to very strong
Husk (presence after harvest)	husked
Ventral furrow hairiness	present
Shape of lemma base	bevelled

Plant

Growth habit	intermediate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	horizontal
Time of ear emergence	medium
Flag leaf glaucosity	strong to very strong
Ear glaucosity	strong
Ear attitude (21 days after ear emergence)	semi-erect
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	weak
Awn tips	medium to strong
Lemma nerves (at early dough)	weak to medium

Ear

Number of rows	six
Shape	parallel
Density	medium to dense
Length (excluding awns)	short
Awn length (compared to ear)	long
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	n/a
Attitude	n/a

Rachis

Length of first segment	short
Curvature of first segment	very weak

JORDAN**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	very lax to lax
Length	medium to long
Shape	tapering

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	medium to long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	weak to medium
Ear - glaucosity	weak to medium
Ear - attitude	semi-erect to horizontal

CHARACTERS IN DETAIL**Grain**

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	absent or very weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	present
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	semi-erect
Time of ear emergence	medium to late
Flag leaf glaucosity	strong
Ear glaucosity	very weak to weak
Ear attitude (21 days after ear emergence)	semi-erect
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	absent or very weak
Awn tips	absent or very weak
Lemma nerves (at early dough)	absent or very weak

Ear

Number of rows	two
Shape	slightly tapering
Density	medium
Length (excluding awns)	medium
Awn length (compared to ear)	medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	not applicable

Rachis

Length of first segment	medium
Curvature of first segment	weak to medium

Variety Name: **KWS Astaire**

Crop:

Breeders Reference:

AFP No: 2/2877

Breeder: KWS UK Ltd

Translated from: German

Origin:

Date: 23 March 2017

Code: KAST

*Plant - growth habit:	Intermediate to semi prostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Medium
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Strong to very strong
*Ear- glaucosity:	Medium to strong
Ear- attitude:	Semi erect
Ear- length:	Medium
*Ear- number of rows:	More than two
*Ear- density:	Medium
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Medium to long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn cf grain:	Very long
Lemma nerves- anthocyanin colouration:	
*Plant height:	Medium
Collar type:	
Rachis- length of first segment:	Medium to long
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium to strong
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Strongly coloured
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Medium
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

28 March 2017

KWS BIRDIS**CHARACTERS IN DETAIL****Grain**

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	strong
Husk (presence after harvest)	husked
Ventral furrow hairiness	present
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	horizontal
Time of ear emergence	medium
Flag leaf glaucosity	strong
Ear glaucosity	strong
Ear attitude (21 days after ear emergence)	erect to semi-erect
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	medium
Awn tips	medium
Lemma nerves (at early dough)	very weak to weak

Ear

Number of rows	six
Shape	parallel
Density	medium to dense
Length (excluding awns)	very short to short
Awn length (compared to ear)	long to very long
Median spikelet - length of glume+awn cf grain	slightly longer

Sterile Spikelet

Development	not applicable
Attitude	not applicable

Rachis

Length of first segment	short to medium
Curvature of first segment	medium to strong

KWS CARBIS**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	very short to short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect

KWS CARIBOU**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax to medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short
Lower leaf - hairiness of sheaths	absent
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude	semi-erect

KWS CASSIA**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium
Shape	parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	short
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	very weak to weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

KWS FEERIS**CHARACTERS IN DETAIL****Ear**

Number of rows	more than two
Density	dense
Length	short to medium
Shape	parallel

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	short
Curvature of first segment	weak

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn cf grain	longer

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	medium
Lemma nerves - at early dough	weak
Awn tips	strong

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	strong to very strong
Ear - glaucosity	strong
Ear - attitude	erect to semi-erect

KWS GIMLET**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	very lax to lax
Length	long to very long
Shape	tapering

Awn

Length compared to ear	short to medium
------------------------	-----------------

Rachis

Length of first segment	long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	medium
Lower leaf - hairiness of sheaths	absent
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude	horizontal

KWS GLACIER**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	erect to semi-erect

KWS HAWKING**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	absent
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	weak to medium
Ear - attitude	semi-erect

KWS HERACLIS**CHARACTERS IN DETAIL****Grain**

Aleurone colour	whitish
Rachilla hair type	short
Spiculation of inner lateral nerves	weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	present
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	semi-erect to horizontal
Time of ear emergence	medium to late
Flag leaf glaucosity	medium
Ear glaucosity	weak
Ear attitude (21 days after ear emergence)	semi-erect to horizontal
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	absent or very weak
Awn tips	absent or very weak
Lemma nerves (at early dough)	absent or very weak

Ear

Number of rows	two
Shape	slightly tapering
Density	sparse to medium
Length (excluding awns)	medium to long
Awn length (compared to ear)	medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	n/a

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

KWS MERIDIAN**CHARACTERS IN DETAIL****Ear**

Number of rows	six
Density	medium
Length	short to medium
Shape	parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	n/a
Median spikelet - length of glume+awn cf grain	longer

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	absent to very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent to very weak

Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium to strong
Ear - attitude 21 days after ear emerg.	semi-erect

KWS ORWELL**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	long
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate
Plant - length	very short to short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	weak
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	erect to semi-erect

KWS TARDIS**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax to medium
Length	long
Shape	tapering

Awn

Length compared to ear	short to medium
------------------------	-----------------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	short
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium
Ear - attitude	semi-erect

KWS TOWER**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	long
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	erect to semi-erect

KWS VALENCIS**CHARACTERS IN DETAIL****Grain**

Aleurone colour	whitish
Rachilla hair type	short
Spiculation of inner lateral nerves	weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	absent
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	semi-erect
Time of ear emergence	medium to late
Flag leaf glaucosity	medium
Ear glaucosity	weak
Ear attitude (21 days after ear emergence)	semi-erect
Length (stem, ears and awns)	medium to long

Anthocyanin colouration

Flag leaf auricles	absent or very weak
Awn tips	absent or very weak
Lemma nerves (at early dough)	absent or very weak

Ear

Number of rows	two
Shape	slightly tapering
Density	medium
Length (excluding awns)	long
Awn length (compared to ear)	short
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	n/a

Rachis

Length of first segment	long to very long
Curvature of first segment	weak to medium

LG CAMPUS**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	short
------------------------	-------

Rachis

Length of first segment	short to medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	husked
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	very weak to weak
Ear - attitude	semi-erect to horizontal

LG CAPITOL**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium to dense
Length	medium to long
Shape	tapering

Awn

Length compared to ear	short to medium
------------------------	-----------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	husked
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude	semi-erect

Variety Name: **LG CAPRICORN**

Crop: **Winter barley**

Breeders Reference:

AFP No: 2/2735

Breeder:

Translated from: French

Origin:

Date: 01 March 2016

Code: LCORN

*Plant - growth habit:	Intermediate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Early
*Flag leaf- anthocyanin colouration of auricles:	Absent to very weak
Frequency of plants with recurved flag leaves:	Medium
Flag leaf- glaucosity of sheath:	Strong
*Ear- glaucosity:	Strong
Ear- attitude:	Semi erect
Ear- length:	Medium
*Ear- number of rows:	Two
*Ear- density:	Lax to medium
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Absent to very weak
*Awn- length compared to ear:	Medium to long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	Deficiens
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn cf grain:	
Lemma nerves- anthocyanin colouration:	Absent to very weak
*Plant height:	Medium
Collar type:	
Rachis- length of first segment:	
Rachis- width of first segment:	
Rachis- curvature of first segment:	
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Absent to very weak
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent to very weak
*Grain- ventral furrow - presence of hairs:	Present
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Lateral
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

07 April 2025

LG CARAVELLE**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium to dense
Length	short to medium
Shape	tapering

Awn

Length compared to ear	medium to long
------------------------	----------------

Rachis

Length of first segment	short to medium
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	husked
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	absent
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium to strong
Ear - attitude	semi-erect

LG CARPENTER**CHARACTERS IN DETAIL****Grain**

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	very weak to weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	absent
Shape of lemma base	bevelled

Plant

Growth habit	intermediate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	horizontal
Time of ear emergence	medium to late
Flag leaf glaucosity	medium
Ear glaucosity	weak to medium
Ear attitude (21 days after ear emergence)	horizontal
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	absent or very weak
Awn tips	absent or very weak
Lemma nerves (at early dough)	absent or very weak

Ear

Number of rows	two
Shape	slightly tapering
Density	medium
Length (excluding awns)	medium to long
Awn length (compared to ear)	short
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	n/a

Rachis

Length of first segment	long to very long
Curvature of first segment	weak to medium

LG CATAPULT

CHARACTERS IN DETAIL

Grain

Aleurone colour	dark grey blue
Rachilla hair type	long
Spiculation of inner lateral nerves	absent or very weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	absent
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	horizontal
Time of ear emergence	late
Flag leaf glaucosity	strong
Ear glaucosity	medium to strong
Ear attitude (21 days after ear emergence)	horizontal to semi-drooping
Length (stem, ears and awns)	short to medium

Anthocyanin colouration

Flag leaf auricles	absent or very weak
Awn tips	absent or very weak
Lemma nerves (at early dough)	absent or very weak

Ear

Number of rows	two
Shape	slightly tapering
Density	medium
Length (excluding awns)	medium to long
Awn length (compared to ear)	short to medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	not applicable

Rachis

Length of first segment	medium to long
Curvature of first segment	medium

LG DAZZLE**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	long
Shape	tapering

Awn

Length compared to ear	short
------------------------	-------

Rachis

Length of first segment	medium to long
Curvature of first segment	strong to very strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	strong
Ear - glaucosity	medium
Ear - attitude	semi-erect to horizontal

LG FLYNN**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

Awn

Length compared to ear	short to medium
------------------------	-----------------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	medium to strong
Lemma nerves - at early dough	medium to strong
Awn tips	strong

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude	semi-erect

LG MOUNTAIN**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	short to medium
------------------------	-----------------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	semi-erect to intermediate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude	erect to semi-erect

LIGHTNING**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	very lax to lax
Length	long
Shape	tapering

Awn

Length compared to ear	short
------------------------	-------

Rachis

Length of first segment	medium
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	weak to medium
Ear - attitude	semi-erect to horizontal

MARIS OTTER**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium to long
Shape	parallel

Awn

Length compared to ear	long (longer than ear)
------------------------	------------------------

Rachis

Length of first segment	medium to long
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	absent/v. weak (0-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	medium
Awn tips	medium

Plant growth characters

Plant - growth habit	semi-prostrate to prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late to very late
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	horizontal

Variety Name: **Memento**

Crop: Winter Barley

Breeders Reference: SC 16666 OH

AFP No:

Breeder: Secobra Recherches

Translated from: French

Origin:

Date: 04 April 2019

Code: Meme

*Plant - growth habit:	Intermediate
*Lower leaves - hairiness of leaf sheaths:	
*Time of ear emergence:	Early to medium
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	
*Ear- glaucosity:	Medium to strong
Ear- attitude:	Horizontal
Ear- length:	Medium
*Ear- number of rows:	Two
*Ear- density:	Dense
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	NA
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn cf grain:	
Lemma nerves- anthocyanin colouration:	Absent or very weak
*Plant height:	Medium
Collar type:	
Rachis- length of first segment:	Medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Weak to medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Strongly coloured
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Weak to medium
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Absent or very weak
*Grain- ventral furrow - presence of hairs:	Present
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

11 April 2019

NOS OLENA

CHARACTERS IN DETAIL

Grain

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	absent or very weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	present
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	semi-erect
Time of ear emergence	medium to late
Flag leaf glaucosity	medium to strong
Ear glaucosity	weak
Ear attitude (21 days after ear emergence)	semi-erect
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	weak
Awn tips	weak
Lemma nerves (at early dough)	medium to strong

Ear

Number of rows	two
Shape	slightly tapering
Density	medium
Length (excluding awns)	medium to long
Awn length (compared to ear)	medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	not applicable

Rachis

Length of first segment	long
Curvature of first segment	medium to strong

ORGANA

CHARACTERS IN DETAIL

Grain

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	absent or very weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	absent
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	semi-erect
Time of ear emergence	medium to late
Flag leaf glaucosity	medium to strong
Ear glaucosity	weak to medium
Ear attitude (21 days after ear emergence)	semi-erect
Length (stem, ears and awns)	long

Anthocyanin colouration

Flag leaf auricles	absent or very weak
Awn tips	absent or very weak
Lemma nerves (at early dough)	absent or very weak

Ear

Number of rows	two
Shape	slightly tapering
Density	medium
Length (excluding awns)	medium
Awn length (compared to ear)	medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	not applicable

Rachis

Length of first segment	medium to long
Curvature of first segment	medium

PEARL**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	long (longer than ear)
------------------------	------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	divergent
Sterile spikelet - development	
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	medium (3-5 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	medium to strong
Lemma nerves - at early dough	very weak to weak
Awn tips	weak to medium

Plant growth characters

Plant - growth habit	intermediate
Plant - length	medium to long
Lower leaf - hairiness of sheaths	present
Time of ear emergence	late
Flag leaf - glaucosity of sheath	strong to very strong
Ear - glaucosity	strong
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

RESOLUTE**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	sparse to medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	long
Curvature of first segment	weak to medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	husked
Spiculation of inner lateral nerves	absent or v. weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	weak to medium
Ear - glaucosity	weak
Ear - attitude	erect to semi-erect

RGT ORIENTATION

CHARACTERS IN DETAIL

Grain

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	absent or very weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	absent
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	semi-erect to horizontal
Time of ear emergence	medium to late
Flag leaf glaucosity	medium to strong
Ear glaucosity	weak to medium
Ear attitude (21 days after ear emergence)	semi-erect to horizontal
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	absent or very weak
Awn tips	absent or very weak
Lemma nerves (at early dough)	absent or very weak

Ear

Number of rows	two
Shape	slightly tapering
Density	medium
Length (excluding awns)	medium to long
Awn length (compared to ear)	medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	not applicable

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

ROSEMARY**CHARACTERS IN DETAIL****Grain**

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	medium to strong
Husk (presence after harvest)	husked
Ventral furrow hairiness	absent
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	horizontal
Time of ear emergence	medium to late
Flag leaf glaucosity	medium
Ear glaucosity	weak
Ear attitude (21 days after ear emergence)	semi-erect to horizontal
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	absent or very weak
Awn tips	absent or very weak
Lemma nerves (at early dough)	absent or very weak

Ear

Number of rows	two
Shape	slightly tapering
Density	medium
Length (excluding awns)	medium to long
Awn length (compared to ear)	medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	not applicable

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

CHARACTERS IN DETAIL**Grain**

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	absent or very weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	present
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	horizontal
Time of ear emergence	medium to late
Flag leaf glaucosity	weak to medium
Ear glaucosity	weak
Ear attitude (21 days after ear emergence)	semi-erect to horizontal
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	medium to strong
Awn tips	strong
Lemma nerves (at early dough)	strong to very strong

Ear

Number of rows	two
Shape	slightly tapering
Density	sparse
Length (excluding awns)	long to very long
Awn length (compared to ear)	medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	n/a

Rachis

Length of first segment	medium
Curvature of first segment	medium

SAFFRON**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	medium
Length	medium
Shape	parallel

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	short to medium
Curvature of first segment	medium

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	strongly coloured ("blue")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak to medium
Rachilla hair type	long
Ventral furrow - presence of hairs	present
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	
Lemma nerves - at early dough	absent or very weak
Awn tips	

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	absent or very weak
Ear - attitude 21 days after ear emerg.	semi-erect to horizontal

Variety Name: **Sensation**

Crop:

Breeders Reference: LEU73203

AFP No:

Breeder: Deutssche Saatveredelung

Translated from:

Origin:

Date: 27 May 2020

Code: B3295

*Plant - growth habit:	Intermediate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Very early to early
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Strong
*Ear- glaucosity:	Medium
Ear- attitude:	Horizontal
Ear- length:	Medium to long
*Ear- number of rows:	More than two
*Ear- density:	Medium
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	
*Awn- length compared to ear:	Long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	NA
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn cf grain:	Equal
Lemma nerves- anthocyanin colouration:	
*Plant height:	
Collar type:	
Rachis- length of first segment:	Medium to long
Rachis- width of first segment:	
Rachis- curvature of first segment:	Weak to medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Short
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Strong
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Weak
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Alternative
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

04 May 2022

CHARACTERS IN DETAIL**Grain**

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	strong to very strong
Husk (presence after harvest)	husked
Ventral furrow hairiness	present
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	semi-erect
Time of ear emergence	medium to late
Flag leaf glaucosity	medium to strong
Ear glaucosity	weak to medium
Ear attitude (21 days after ear emergence)	semi-erect to horizontal
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	weak
Awn tips	medium to strong
Lemma nerves (at early dough)	medium

Ear

Number of rows	six
Shape	parallel
Density	medium to dense
Length (excluding awns)	medium
Awn length (compared to ear)	medium
Median spikelet - length of glume+awn cf grain	much longer

Sterile Spikelet

Development	n/a
Attitude	n/a

Rachis

Length of first segment	medium
Curvature of first segment	weak to medium

SU ARION**CHARACTERS IN DETAIL****Grain**

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	very weak to weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	absent
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	semi-erect
Time of ear emergence	medium to late
Flag leaf glaucosity	strong
Ear glaucosity	very weak to weak
Ear attitude (21 days after ear emergence)	semi-erect to horizontal
Length (stem, ears and awns)	medium to long

Anthocyanin colouration

Flag leaf auricles	medium to strong
Awn tips	strong
Lemma nerves (at early dough)	strong

Ear

Number of rows	two
Shape	slightly tapering
Density	medium
Length (excluding awns)	medium
Awn length (compared to ear)	medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	n/a

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

SU NEWMARKET**CHARACTERS IN DETAIL****Grain**

Aleurone colour	whitish
Rachilla hair type	short
Spiculation of inner lateral nerves	absent or very weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	absent
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	present
Flag leaf attitude	semi-erect
Time of ear emergence	medium
Flag leaf glaucosity	strong
Ear glaucosity	medium
Ear attitude (21 days after ear emergence)	semi-erect
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	weak
Awn tips	weak
Lemma nerves (at early dough)	weak

Ear

Number of rows	two
Shape	strongly tapering
Density	sparse to medium
Length (excluding awns)	medium to long
Awn length (compared to ear)	very short to short
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	n/a

Rachis

Length of first segment	medium to long
Curvature of first segment	weak to medium

SURGE**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	short to medium
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	weakly coloured
Husk - presence after harvest	present
Spiculation of inner lateral nerves	v. weak to weak
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	absent or very weak
Lemma nerves - at early dough	absent or very weak
Awn tips	absent or very weak

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	very short to short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	weak to medium
Ear - glaucosity	weak
Ear - attitude 21 days after ear emerg.	semi-erect

SY VENTURE

CHARACTERS IN DETAIL

Ear

Number of rows	two
Density	medium
Length	medium
Shape	tapering

Awn

Length compared to ear	medium (+/- equal to ear)
------------------------	---------------------------

Rachis

Length of first segment	medium
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong (5-10 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	very strong
Awn tips	strong

Plant growth characters

Plant - growth habit	intermediate to semi-prostrate
Plant - length	short
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium
Ear - glaucosity	medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

SY VESSEL**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax to medium
Length	medium to long
Shape	tapering

Awn

Length compared to ear	medium
------------------------	--------

Rachis

Length of first segment	medium
Curvature of first segment	strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish
Husk - presence after harvest	present
Spiculation of inner lateral nerves	strong to v. strong
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping

Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	strong to very strong
Awn tips	strong

Plant growth characters

Plant - growth habit	semi-prostrate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	medium
Ear - attitude	semi-erect

TALISMAN**CHARACTERS IN DETAIL****Ear**

Number of rows	two
Density	lax
Length	medium to long
Shape	tapering

Awn

Length compared to ear	short (shorter than ear)
------------------------	--------------------------

Rachis

Length of first segment	medium to long
Curvature of first segment	medium to strong

Spikelet

Sterile spikelet - attitude	n/a
Sterile spikelet - development	none or rudimentary(deficiens)
Median spikelet - length of glume+awn cf grain	equal

Grain

Kernel - colour of aleurone layer	whitish ("white")
Husk - presence after harvest	present
Spiculation of inner lateral nerves	weak (1-2 per nerve)
Rachilla hair type	long
Ventral furrow - presence of hairs	absent
Disposition of lodicules	clasping ("collar" type)

Anthocyanin colouration

Flag leaf - auricles	strong
Lemma nerves - at early dough	strong
Awn tips	strong

Plant growth characters

Plant - growth habit	intermediate
Plant - length	short to medium
Lower leaf - hairiness of sheaths	present
Time of ear emergence	medium to late
Flag leaf - glaucosity of sheath	medium to strong
Ear - glaucosity	weak to medium
Ear - attitude 21 days after ear emerg.	erect to semi-erect

Variety Name: Valerie

Crop:

Breeders Reference: BR 11500 R 6

AFP No:

Breeder: Saatzeit Josef Breun

Translated from: German

Origin:

Date: 04 April 2019

Code: Valer

*Plant - growth habit:	Intermediate to semi-prostrate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Early to medium
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Strong to very strong
*Ear- glaucosity:	Medium
Ear- attitude:	Semi recurved
Ear- length:	Medium to long
*Ear- number of rows:	Two
*Ear- density:	Medium
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	Medium
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	Parallel
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn cf grain:	
Lemma nerves- anthocyanin colouration:	Equal
*Plant height:	Short to medium
Collar type:	
Rachis- length of first segment:	Medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium to strong
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Absent or very weak
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	
*Grain- ventral furrow - presence of hairs:	
*Grain- husk:	
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	
*Seasonal type:	
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

11 April 2019

Variety Name: **Valiant**

Crop: Winter Barley

Breeders Reference: AC 14/361/28

AFP No:

Breeder: Elsoms Ackermann Barley Ltd

Translated from:

Origin:

Date: 28 March 2024

Code: VALI

*Plant - growth habit:	Intermediate
*Lower leaves - hairiness of leaf sheaths:	Present
*Time of ear emergence:	Medium
*Flag leaf- anthocyanin colouration of auricles:	Absent or very weak
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Medium to strong
*Ear- glaucosity:	Weak to medium
Ear- attitude:	Horizontal
Ear- length:	Long
*Ear- number of rows:	Two
*Ear- density:	Medium
Ear- shape:	Slightly tapering
*Awn- anthocyanin colouration of the tips:	Absent or very weak
*Awn- length compared to ear:	
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	NA
Sterile spikelet-shape of tip:	NA
Sterile spikelet- length of lemma:	NA
Median spikelet- length of glume + awn cf grain:	Equal
Lemma nerves- anthocyanin colouration:	Absent or very weak
*Plant height:	Short
Collar type:	
Rachis- length of first segment:	Medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium to strong
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Very weak to weak
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Husked
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Bevelled
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

09 April 2024

VITALITY**CHARACTERS IN DETAIL****Grain**

Aleurone colour	whitish
Rachilla hair type	long
Spiculation of inner lateral nerves	absent or very weak
Husk (presence after harvest)	husked
Ventral furrow hairiness	present
Shape of lemma base	bevelled

Plant

Growth habit	intermediate to semi-prostrate
Green colour (at tillering)	medium
Lower leaf sheath hairiness	absent
Flag leaf attitude	horizontal
Time of ear emergence	medium to late
Flag leaf glaucosity	medium
Ear glaucosity	medium
Ear attitude (21 days after ear emergence)	semi-erect to horizontal
Length (stem, ears and awns)	medium

Anthocyanin colouration

Flag leaf auricles	absent or very weak
Awn tips	absent or very weak
Lemma nerves (at early dough)	absent or very weak

Ear

Number of rows	two
Shape	slightly tapering
Density	sparse to medium
Length (excluding awns)	long to very long
Awn length (compared to ear)	medium
Median spikelet - length of glume+awn cf grain	equal

Sterile Spikelet

Development	none or rudimentary
Attitude	not applicable

Rachis

Length of first segment	long
Curvature of first segment	strong

Variety Name: **Zophia**

Crop: Winter Barley

Breeders Reference: UN 116261

AFP No:

Breeder: Sejet

Translated from: French

Origin:

Date: 04 April 2019

Code: ZOPH

*Plant - growth habit:	Intermediate
*Lower leaves - hairiness of leaf sheaths:	
*Time of ear emergence:	Early
*Flag leaf- anthocyanin colouration of auricles:	Medium to strong
Frequency of plants with recurved flag leaves:	
Flag leaf- glaucosity of sheath:	Strong
*Ear- glaucosity:	Medium to strong
Ear- attitude:	Horizontal
Ear- length:	
*Ear- number of rows:	Two
*Ear- density:	Medium to dense
Ear- shape:	Parallel
*Awn- anthocyanin colouration of the tips:	Medium
*Awn- length compared to ear:	Long
Awn- spiculation of margins:	
*Sterile spikelet-attitude:	
Sterile spikelet-shape of tip:	
Sterile spikelet- length of lemma:	
Median spikelet- length of glume + awn cf grain:	Equal
Lemma nerves- anthocyanin colouration:	Medium
*Plant height:	
Collar type:	
Rachis- length of first segment:	Medium
Rachis- width of first segment:	
Rachis- curvature of first segment:	Medium
Rachis- shape of first segment:	
Rachis(2-row)- humping of segments (mid third):	
Rachis(6-row)- degree of zig-zag (mid third):	
Kernel- colour of aleurone layer:	Whitish
*Grain- rachilla hair type:	Long
Grain- rachilla length:	
Grain- rachilla - length of tip hairs:	
Grain- spiculation of inner lateral nerves:	Strong
Grain- spiculation of outer lateral nerves:	
Grain- anthocyanin colouration of lemma nerves:	Medium
*Grain- ventral furrow - presence of hairs:	Absent
*Grain- husk:	Present
Grain- length of lodicules ("collar" type only):	
Grain- disposition of lodicules:	Clasping
*Seasonal type:	Winter
Extra characters:	

* = Characteristics which should always be included in the description of the variety.

11 April 2019