Variety Name: **Aarden** crop: Winter Wheat

Breeder's Reference: SUR 97271/10 AFP No: 1/1573

Breeder: Saaten Union Recherche SARL Translated from: German

Origin: Date: 19 November 2003

Code: ARDN

Coleoptile-anthocyanin colouration: medium

*Plant - growth habit: semi-prostrate to prostrate

*Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: absent or very low

*Flag leaf - Glaucosity of sheath: strong
Flag leaf - Glaucosity of leaf blade: strong

Anthers - anthocyanin colouration:

Glaucosity of neck: strong to very strong

Hairiness of uppermost node:

*Plant - height: short

*Straw - wall thickness @ 20cms below ear: very thin

*Ear - Glaucosity: strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: white

*Ear - Shape (side profile): tapering
Ear - Length: medium

*Ear - Density: medium

*Awns or scurs - presence: scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: short

*Awns at tip of ear - length: Lower glume - Beak length:

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width: Lower glume - Shoulder type:

Lower glume - External surface roughness: Lower glume - Extent of internal hairs:

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour: red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: medium *Seasonal type: winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 06 May 2020

AEROBIC

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: dense

Shape (side profile) parallel sided
Awns or scurs - presence: awns present

Awns or scurs - distribution:

Awns or Scurs - length: very long
Apical rachis segment - convex face hairiness: weak

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: weak to medium - group 1 to 2

Beak length: medium to long
Beak shape (curvature): slightly curved

Shoulder narrow

Shoulder slightly sloping

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: medium

Straw

Length: short
Wall thickness @ 20cms below ear: very thick

Glaucosity

Flag leaf - sheath: weak to medium
Flag leaf - blade (lower side): very weak to weak

Culm neck: weak

Ear - Glaucosity: absent or very weak

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: semi-prostrate

Time ear emergence: early

Flag leaf - Anthocyanin colouration of auricles:



ALCHEMY

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼
Awns or scurs - length: short

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



ALMARA

long

CHARACTERS IN DETAIL

Awns or scurs - length

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4

Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width medium

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-prostrate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



AMBOISE

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium



AMBROSIA

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)
Beak length medium to long
Beak curvature moderately curved

Shoulder width narrow Shoulder shape straight

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



AMPLIFY

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering

Awns or scurs - presence scurs present

Awns or scurs - distribution upper 1/2

Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong (2+)
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



Variety Name: ANAPOLIS crop: Winter wheat

Breeder's Reference: AFP No: 1/2148

Breeder: NORDSAAT Translated from: German

Origin: Date: 23 July 2013

Code: ANAP

Coleoptile-anthocyanin colouration:Medium to strong*Plant - growth habit:Semi prostrate*Time ear emergence:Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:*Flag leaf - Glaucosity of sheath:

Very low to low

Strong to very strong

Flag leaf - Glaucosity of leaf blade: Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong to very strong

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Absent or very thin.

*Ear - Glaucosity: Strong to very strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Tapering

Ear - Length: Medium

*Ear - Density: Lax to medium

*Awns or scurs - presence: Present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short to short

*Awns at tip of ear - length:

Lower glume - Beak length:Very short to shortLower glume - Beak shape (curvature):Slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width:NarrowLower glume - Shoulder type:Straight

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Medium

Lower glume - Internal imprint:

Lower lemma - Beak shape: Slightly curved

Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour:

Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: Medium to dark

*Seasonal type: Winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 31 March 2015

ARARAT

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: lax

Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short

Apical rachis segment - convex face hairiness: very weak to weak

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: medium - group 2

Beak length: medium

Beak shape (curvature): moderately curved

Shoulder medium
Shoulder elevated

Lower Lemma

Beak shape: slightly curved

Grain

Colour red
Colouration with phenol: medium

Straw

Length: medium Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: medium to strong
Flag leaf - blade (lower side): medium to strong

Culm neck: strong

Ear - Glaucosity: strong to very strong

Plant growth characters

Coleoptile-anthocyanin medium

Plant - growth habit: intermediate to semi-prostrate

Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



Variety Name: Arkeos

Crop:

Breeder's Reference: AFP No:

Breeder: Nickerson International Research **Translated from:**

Origin: Date: 26 January 2015

Code: ARKE

Coleoptile-anthocyanin colouration: Medium

*Plant - growth habit: Semi-prostrate to prostrate

*Time ear emergence: Early

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:

*Flag leaf - Glaucosity of sheath: Strong to very strong Flag leaf - Glaucosity of leaf blade: Strong to very strong

Anthers - anthocyanin colouration:

Glaucosity of neck:

Hairiness of uppermost node:

*Plant - height: Short to medium *Straw - wall thickness @ 20cms below ear: Very thin

Medium *Ear - Glaucosity:

Ear - Supernumerary spikelets - frequency:

White *Ear - Colour:

*Ear - Shape (side profile):

Ear - Length: Long

*Ear - Density: Lax to medium *Awns or scurs - presence: Present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Short to medium

*Awns at tip of ear - length:

Lower glume - Beak length: Short to medium

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow

Lower glume - Shoulder type: Straight to slightly sloping

Lower glume - External surface roughness: Lower glume - Extent of internal hairs:

Lower glume - Internal imprint:

Medium to strong Lower lemma - Beak shape: Slightly curved Lower lemma - Beak length-: Short to medium

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Medium *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: Mdium to dark Winter *Seasonal type:

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 30 March 2015

ASTOUND

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long

Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4

Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs medium to strong (group 2+)

Acette of international and a second for the second

Beak length long

Beak curvature strongly curved Shoulder width medium

Shoulder shape elev. to strongly elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong

Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



AVATAR

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density dense Shape-side profile parallel sided

Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4

Awns or scurs - length: medium to long weak

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak Plant - growth habit semi-erect Time of ear emergence medium to late



Variety Name: **Axioma** Crop: Wheat

Breeder's Reference: SEC 228-02-4 AFP No:

Breeder: Secobra Rechereches Translated from: German

Origin: Date: 19 November 2018

Code: Axio

Coleoptile-anthocyanin colouration:Medium*Plant - growth habit:Intermediate*Time ear emergence:Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Very low to low

*Flag leaf - Glaucosity of sheath: Srong
Flag leaf - Glaucosity of leaf blade: Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Medium *Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour:

*Ear - Shape (side profile): Tapering

Ear - Length: Short to medium

*Ear - Density: Dense

*Awns or scurs - presence: Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short to short

*Awns at tip of ear - length:

Lower glume - Beak length: Short

Lower glume - Beak shape (curvature): Moderately curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow to medium

Lower glume - Shoulder type: Straight

Lower glume - External surface roughness: Lower glume - Extent of internal hairs: Lower glume - Internal imprint:

Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Weak *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: *Seasonal type:Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 20 March 2019



BAHAMAS CS

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium
Beak curvature slightly curved
Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



BAMFORD

CHARACTERS IN DETAIL

Ear

Colour white Length medium to long Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish
Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



BANJO

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol dark to very dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong to very strong
Plant - growth habit semi-prostrate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



BANQUO

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



BARKSTON

long to very long

CHARACTERS IN DETAIL

Awns or scurs - length

Ear

Colour white

medium to long Length Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2

Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong (group 3) Beak length medium Beak curvature slightly curved Shoulder width narrow elevated

Shoulder shape

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) medium to strong

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermed. to semi-prostrate

Time of ear emergence very late



BATSMAN

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermediate
Time of ear emergence late to very late



BATTALION

CHARACTERS IN DETAIL

Ear

Colour white

Length medium

Density lax to medium

Shape-side profile tapering

Awns or scurs - presence scurs present

Awns of scurs - distribution upper ¼

Awns or scurs - length: medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium - Group 2
Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence early to medium



BELENUS

CHARACTERS IN DETAIL

Ear

Colour: coloured

Length: short

Density: medium

Shape (side profile) parallel sided

Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: medium
Apical rachis segment - convex face hairiness: weak

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: weak - group 1
Beak length: very short

Beak shape (curvature): slightly curved

Shoulder narrow

Shoulder slightly sloping

Lower Lemma

Beak shape: slightly curved

Grain

Colour red
Colouration with phenol: medium

Straw

Length: short Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: weak

Flag leaf - blade (lower side): absent or very weak

Culm neck: absent or very weak

Ear - Glaucosity: absent or very weak

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: intermediate Time ear emergence: medium to late Flag leaf - Anthocyanin colouration of auricles: absent or very weak



BELEPI

CHARACTERS IN DETAIL

Ear

Colour: white

Length: long to very long

Density: medium
Shape (side profile) parallel

Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: medium
Apical rachis segment - convex face hairiness: weak

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3

Beak length:

Beak shape (curvature):

Shoulder straight

Shoulder

Lower Lemma

Beak shape: straight to slightly curved

Grain

Colour red

Colouration with phenol: dark to very dark

Straw

Length: medium Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong to very strong
Flag leaf - blade (lower side): strong to very strong

Culm neck:strong to very strongEar - Glaucosity:strong to very strong

Plant growth characters

Coleoptile-anthocyanin weak

Plant - growth habit: intermediate to semi-prostrate

Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:



BELGRADE

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surfaceroughExtent of internal hairsstrong (3)Beak lengthmedium to longBeak curvaturemoderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



BELUGA

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: short to medium

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit semi-erect
Time of ear emergence medium



BENNINGTON

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlongApical rachis segment - convex face hairinessstrong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Flag leaf - anthocyanin of auricles

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermediate
Time of ear emergence very late



BISCAY

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax Shape-side profile tapering Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-prostrate

Time of ear emergence late
Flag leaf - anthocyanin of auricles very weak



BLACKSMITH

CHARACTERS IN DETAIL

Ear

Colour white

medium to long Length Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness absent or very small

Anthers - anthocyanin colouration present

Lower glume

External surface smooth Extent of internal hairs weak (group 1) Beak length short to medium Beak curvature slightly curved

Shoulder width very narrow to narrow Shoulder shape slightly elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour purple very dark

Colouration with phenol

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very strong Plant - growth habit intermediate Time of ear emergence medium to late Flag leaf - anthocyanin of auricles absent or very weak



BLACKSTONE

CHARACTERS IN DETAIL

Ear

white Colour medium Length Density dense Shape-side profile parallel sided Awns or scurs - presence scurs present upper 3/4 Awns or scurs - distribution Awns or scurs - length medium to long Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium Shoulder shape slightly elevated

Lower lemma

Beak curvature geniculate

Grain

Colour reddish
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) medium to strong
Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



BOLINDER

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs weak to medium (group 1+)

Beak length short to medium

Beak curvature moderate to strongly curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour reddish
Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



BOOMER

CHARACTERS IN DETAIL

Ear

Colour: white
Length: medium
Density: medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short
Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong -group 3

Beak length: short

Beak shape (curvature): moderately curved

Shoulder narrow
Shoulder elevated

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: dark

Straw

Length: short Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): strong
Culm neck: strong

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin strong

Plant - growth habit: semi-prostrate
Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:



BRIDGEHEAD

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late to very late



BRITANNIA

absent

CHARACTERS IN DETAIL

Anthers - anthocyanin colouration

Ear

Colour white

Length medium to long

Density lax

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length very long
Apical rachis segment - convex face hairiness strong

Lower glume

External surface slightly rough
Extent of internal hairs weak to medium (1+)

Beak length medium to long
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong

Culm - neck medium

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late to very late



BROMPTON

CHARACTERS IN DETAIL

Ear

white Colour Length medium lax to medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak lengthshort to mediumBeak curvatureslightly curvedShoulder widthnarrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



BUTLER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length medium
Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length short

Beak curvature slightly curved Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Ear

Flag leaf - sheath absent or very weak
Flag leaf- blade (lower side) absent or very weak
Culm - neck absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

absent or very weak

late

Time of ear emergence



BYSTRA

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium
Density: medium to dense
Shape (side profile) parallel sided
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness: strong

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: weak - group 1
Beak length: very short

Beak shape (curvature): slightly curved

Shoulder narrow

Shoulder slightly sloping

Lower Lemma

Beak shape: slightly curved

Grain

Colour red

Colouration with phenol: light to medium

Straw

Length: short Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): strong
Culm neck: strong

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin weak
Plant - growth habit: semi-erect
Time ear emergence: medium to late
Flag leaf - Anthocyanin colouration of auricles: absent or very weak



CAMBRIDGE

elevated

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¾
Average profile scurs beautiful to the scurs present and several scurs present and

Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature moderately curved
Shoulder width narrow

Lower lemma

Shoulder shape

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium



CASSIUS

CHARACTERS IN DETAIL

Ear

Colour white

Length medium

Density medium

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns of scurs - distribution upper ¼

Awns or scurs - length: long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence late to very late



CATALYST

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



Variety Name: CELLULE crop: Alternative Wheat

Breeder's Reference: fd08114/Cellule AFP No:

Breeder: Florimond Desprez Veuve et Fils **Translated from:**

Origin: Date: 15 May 2018

Code: CELL

Coleoptile-anthocyanin colouration: Very strong

*Plant - growth habit: Intermediate to semi prostrate

*Time ear emergence: Early

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Low

*Flag leaf - Glaucosity of sheath: Very strong

Flag leaf - Glaucosity of leaf blade: Strong ot very strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong to very strong

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Very thin

*Ear - Glaucosity: Strong to very strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Parallel sided
Ear - Length: Medium to long

*Ear - Density: Dense

*Awns or scurs - presence:

Awns or scurs - distibution:

*Scurs at tip of ear - length:

Very long

*Awns at tip of ear - length:

Lower glume - Beak length:Short to mediumLower glume - Beak shape (curvature):Slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width:Narrow ot mediumLower glume - Shoulder type:Slightly sloping or straight

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak to medium

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Weak *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:*Seasonal type:

Alternative

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 16 May 2018

CHAMPION

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length long

Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



CHEVRON

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence early to medium



CHILTON

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributiontip only

Awns or scurs - length: long
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium to long
Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium



CHRONICLE

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium to long
Beak curvature slightly curved

Shoulder width very narrow to narrow Shoulder shape strongly elevated with 2nd pt.

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium



CLAIRE

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: short

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium
Beak curvature slightly curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



COCOON

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs strong - Group 3

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) strong to very strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



COMOROS CS

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



CONQUEROR

CHARACTERS IN DETAIL

Ear

Colour white

Length medium

Density medium

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns of scurs - distribution upper ¼

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



CONSORT

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper ¼

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate
Time of ear emergence medium to late Flag leaf - anthocyanin of auricles absent or very weak



CORDIALE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length short

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect

Time of ear emergence very early to early

Flag leaf - anthocyanin of auricles absent or very weak



CORONATION

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼
Awns or scurs - length: long
Apical rachis segment - convex face hairiness medium

Apical rachis segment - convex face nairiness medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong - Group 2+

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate
Time of ear emergence medium to late



COSTELLO

weak

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs weak to medium (1+)

Beak length short

Beak curvature strongly curved

Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



COUGAR

CHARACTERS IN DETAIL

Ear

Colour white

Length short

Density dense

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns of scurs - distribution upper ½

Awns or scurs - length:

Awns or scurs - length: long to very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



CREEK

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length short

Beak curvature slightly curved Shoulder width narrow Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit semi-erect
Time of ear emergence medium to late



CRUSOE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width medium to broad

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath absent or very weak Flag leaf- blade (lower side) absent or very weak Culm - neck absent or very weak Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



CUBANITA

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: long

Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium to late



DEBEN

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



DELPHI

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper 3/4
Awns or scurs - length: long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit medium to strong semi-erect to intermediate

Time of ear emergence medium to late



DENMAN

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense

Shape-side profiletaperingAwns or scurs - presencescurs presentAwns of scurs - distributionupper ¼Awns or scurs - length:short

Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



DEXTER

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium
Beak length medium
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



DIAMOND

medium to dense

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol medium to dark

Plant

Anthocyanin colouration of coleoptile medium

Growth habit semi-erect to intermediate

Frequency of plants with recurved flag leaves very low to low

Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence medium
Length (stem, ears, awns or scurs) short
Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong

Flag leaf blade (lower side) strong

Ear strong to very strong
Culm - neck strong to very strong

Lower glume

Hairiness of external surface absent

Shoulder width very narrow to narrow Shoulder shape slightly elevated very long

Shape of beak moderate to strongly curved

Extent of internal hairs very large (Group 3)

External surface roughness entire external surface rough

Ear

Density

Length (excluding awns or scurs) short to medium
Awns or scurs - presence awns present
Awns or scurs - length very long

Colour white

Shape in profile parallel sided
Apical rachis hairiness on convex surface small to medium

Awns or scurs - distribution whole length



DICKENS

elevated

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth Extent of internal hairs strong - Group 3 Beak length medium Beak curvature slightly curved Shoulder width narrow Shoulder shape

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) strong to very strong

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong Plant - growth habit semi-erect Time of ear emergence medium



DRUMLIN

CHARACTERS IN DETAIL

Ear

Colour white Length short

Density medium to dense
Shape-side profile tapering

Awns or scurs - presence scurs present

Awns or scurs - distribution upper 1/4

Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



DUNSTON

medium

CHARACTERS IN DETAIL

Ear

white Colour

Length medium to long Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length long

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong (group 3) Beak length medium Beak curvature slightly curved Shoulder width narrow elevated

Shoulder shape

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

short Length

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) medium to strong

Culm - neck medium medium Ear

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence very late



DUXFORD

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow

Shoulder shape strongly elevated with 2nd pt.

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) weak
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit semi-erect
Time of ear emergence early to medium



EDGAR

CHARACTERS IN DETAIL

Ear

Colour: white
Length: long
Density: lax
Shape (side profile) tapering

Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short

Apical rachis segment - convex face hairiness: weak to medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: weak - group 1
Beak length: very short to short
Beak shape (curvature): strongly curved

Shoulder narrow

Shoulder straight to elevated

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: dark

Straw

Length: medium Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): strong

Culm neck: strong to very strong Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin very weak to weak Plant - growth habit: intermediate Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



EDMUNDS

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs strong (3)
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



EINSTEIN

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼
Awns or scurs - length: short

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length short

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck medium to strong
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak
Plant - growth habit semi-erect
Time of ear emergence early to medium
Flag leaf - anthocyanin of auricles absent or very weak



ELABORATE

CHARACTERS IN DETAIL

Ear

Colour white Length long

Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath weak to medium Flag leaf- blade (lower side) weak to medium

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit intermed. to semi-prostrate

late

Time of ear emergence



ELATION

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderately curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

late

Time of ear emergence



ELICIT

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs weak (group 1)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late



ELOQUENCE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Flag leaf - anthocyanin of auricles

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate

Time of ear emergence medium



ELWELL

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium (group 1+)

Beak lengthshort to mediumBeak curvatureslightly curvedShoulder widthnarrow to mediumShoulder shapestraight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence late



ELYSIUM

CHARACTERS IN DETAIL

Ear

white Colour

Length medium to long medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4

Awns or scurs - length long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium Beak curvature slightly curved

Shoulder width very narrow to narrow elev. to strongly elevated Shoulder shape

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red Colouration with phenol dark

Straw

medium Length

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium Plant - growth habit intermediate late

Time of ear emergence



ENERGISE

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium
Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence late



ENERGY

CHARACTERS IN DETAIL

Grain

Colour reddish
Colouration with phenol dark

Plant

Anthocyanin colouration of coleoptile medium to strong
Growth habit intermediate
Frequency of plants with recurved flag leaves
Anthocyanin colouration of flag leaf auricles absent or very low
Time of ear emergence medium to late
Length (stem, ears, awns or scurs) very short to short

Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong
Flag leaf blade (lower side) strong

Ear medium to strong

Culm - neck strong

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium

Shoulder shape slightly elevated to strongly elevated

Length of beak short to medium

Shape of beak moderate to strongly curved

Extent of internal hairs very large (Group 3)

External surface roughness smooth

Ear

Density medium

Length (excluding awns or scurs)short to mediumAwns or scurs - presencescurs presentAwns or scurs - lengthmediumColourwhite

Shape in profile parallel sided

Apical rachis hairiness on convex surface large
Awns or scurs - distribution upper 1/4



EVOLUTION

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper ½

Awns or scurs - length: long to very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak lengthmediumBeak curvaturestrongly curvedShoulder widthmediumShoulder shapeelevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



EXSEPT

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns of scurs - distributiontip onlyAwns or scurs - length:short

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape straight

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-prostrate

Time of ear emergence late to very late

Flag leaf - anthocyanin of auricles absent or very weak



FAIRWAY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution

Awns or scurs - length very short to short
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface
Extent of internal hairs weak to medium (group 1+)

Beak length short

Beak curvature slightly curved Shoulder width narrow Shoulder shape slightly sloping

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-prostrate to prostrate

Time of ear emergence medium to late



FITZROY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs weak to medium (group 1+)

Beak lengthmediumBeak curvaturestrongly curvedShoulder widthmedium to broadShoulder shapestraight to elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong

Culm - neck strong to very strong
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



FLORETT

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium
Density: medium to dense

Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: weak to medium - group 1 to 2

Beak length: very short to short

Beak shape (curvature): moderately curved

Shoulder narrow

Shoulder slightly sloping to straight

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: dark

Straw

Length: short Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong

Flag leaf - blade (lower side): weak to medium

Culm neck: medium to strong

Ear - Glaucosity: medium

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: intermediate Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:



FOREST

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense Shape-side profile parallel sided

Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



FORZOR

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length short

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

smooth External surface Extent of internal hairs medium (2) Beak length short to medium Beak curvature slightly curved Shoulder width narrow to medium elevated

Shoulder shape

Lower lemma

Beak curvature moderately curved

Grain

Colour red Colouration with phenol very dark

Straw

short to medium Length Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium Flag leaf- blade (lower side) weak to medium Culm - neck weak to medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak Plant - growth habit intermediate Time of ear emergence medium to late Flag leaf - anthocyanin of auricles



FREISTON

CHARACTERS IN DETAIL

Ear

Colour white Length long

Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long
Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium to long
Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermed. to semi-prostrate
Time of ear emergence very late



FRENZY

CHARACTERS IN DETAIL

Grain

Colour reddish Colouration with phenol dark

Plant

Anthocyanin colouration of coleoptile strong to very strong Growth habit intermediate

Frequency of plants with recurved flag leaves
Anthocyanin colouration of flag leaf auricles absent or very low absent or very weak Time of ear emergence medium to late

Length (stem, ears, awns or scurs) strong to very strong intermediate

absent or very weak medium to late short

Length (stem, ears, awns or scurs) shor Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong
Flag leaf blade (lower side) strong to very strong
Ear strong to very strong
Culm - neck strong to very strong

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium

Shoulder shape slightly elevated to strongly elevated

Length of beak lor

Shape of beak slight to moderately curved

Extent of internal hairs medium (Group 2)

External surface roughness rough

Ear

Density medium to dense
Length (excluding awns or scurs) medium to long
Awns or scurs - presence scurs present
Awns or scurs - length medium
Colour white
Shape in profile parallel sided

Apical rachis hairiness on convex surface large
Awns or scurs - distribution upper 1/2



Variety Name: Frontal

Crop:

Breeder's Reference: AFP No:

Breeder: Translated from: Dutch

Origin: Date: 24 March 2015

Code: FRNT

Coleoptile-anthocyanin colouration: Absent or very weak

*Plant - growth habit: Intermediate to semi prostrate

*Time ear emergence: Medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:*Flag leaf - Glaucosity of sheath:

Strong to very strong

Flag leaf - Glaucosity of leaf blade: Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong to very strong

Hairiness of uppermost node:

*Plant - height: Medium
*Straw - wall thickness @ 20cms below ear: Very thin

*Ear - Glaucosity: Strong to very strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White
*Ear - Shape (side profile): Tapering

Ear - Length: Short to medium
*Ear - Density: Medium to dense

*Awns or scurs - presence: Present

Awns or scurs - distibution:

*Scurs at tip of ear - length:

*Awns at tip of ear - length: Medium
Lower glume - Beak length: Medium

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width:MediumLower glume - Shoulder type:Elevated

Lower glume - External surface roughness: Lower glume - Extent of internal hairs:

Lower glume - Internal imprint:

Lower lemma - Beak shape: Moderately curved

Lower lemma - Beak length-: Medium

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Medium to strong

*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: Medium to dark

*Seasonal type: Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 30 March 2015



FURLONG

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 3/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (group 3)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



GALLANT

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short - medium
Beak curvature slightly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature straight

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit semi-erect
Time of ear emergence medium



GANDALF

CHARACTERS IN DETAIL

Ear

Colour coloured
Length short to medium

Density medium
Shape-side profile parallel sided
Awns or scurs - presence awns present
Awns or scurs - distribution whole length

Awns or scurs - length very long

Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs medium (group 2)

Beak length very long

Beak curvature moderate to strongly curved
Shoulder width very narrow to narrow
Shoulder shape strongly elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear weak

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-prostrate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles very weak



GARIBALDI

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium (group 2)

Beak length long

Beak curvature strongly curved to geniculate

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



GEFION

CHARACTERS IN DETAIL

Ear

white Colour medium Length Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long Apical rachis segment - convex face hairiness large Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



Variety Name: **Genius** crop: Winter Wheat

Breeder's Reference: AFP No:

Breeder: Nordsaat Saatzuchtgesellschaft Translated from: German

Origin: Date: 08 October 2010

Code: GENI

Coleoptile-anthocyanin colouration:absent or very weak*Plant - growth habit:intermediate*Time ear emergence:medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: very low to low

*Flag leaf - Glaucosity of sheath: strong

Flag leaf - Glaucosity of leaf blade: medium to strong

Anthers - anthocyanin colouration:

Glaucosity of neck: medium to strong

Hairiness of uppermost node:

*Plant - height: short to medium

*Straw - wall thickness @ 20cms below ear: very thin

*Ear - Glaucosity: weak to medium

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: white
*Ear - Shape (side profile): tapering

Ear - Length: medium to long
*Ear - Density: lax to medium
*Awns or scurs - presence: scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: short

*Awns at tip of ear - length:

Lower glume - Beak length: short

Lower glume - Beak shape (curvature): moderately curved

Lower glume - Beak tip:

Lower glume - Shoulder width: narrow

Lower glume - Shoulder type: straight to elevated

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: weak - group 1

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: strong *Grain - colour: red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: medium to dark

*Seasonal type: winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 30 March 2015

GLADIATOR

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface

Extent of internal hairs medium - Group 2

Beak length short

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate

Time of ear emergence early

Flag leaf - anthocyanin of auricles absent or very weak



GLASGOW

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns or scurs - distribution upper 1/4

Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence very early to early Flag leaf - anthocyanin of auricles absent or very weak



GLEAM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4Awns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)

Beak length short to medium

Beak curvature v.slightly curved

Shoulder width very narrow to narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium



GOLDENGUN

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length medium
Beak curvature slightly curved
Shoulder width narrow

Shoulder shape elev. to elev with 2nd point

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong to very strong
Plant - growth habit intermed. to semi-prostrate

Time of ear emergence



GRAFTON

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present

Awns of scurs - distribution upper $\frac{1}{2}$ Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium

Beak curvature moderate to strongly curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium



Variety Name: **Graham** Crop:

Breeder's Reference: AFP No:

Breeder: Syngenta Seeds SAS Translated from: French

Origin: Date: 03 November 2014

Code: GRAH

Coleoptile-anthocyanin colouration: Absent or very weak

*Plant - growth habit: Intermediate to semi prostrate

*Time ear emergence: Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:

*Flag leaf - Glaucosity of sheath: Strong to very strong
Flag leaf - Glaucosity of leaf blade: Strong to very strong

Anthers - anthocyanin colouration:

Glaucosity of neck:

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Very thin

*Ear - Glaucosity: Strong to very strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Parallel sided
Ear - Length: Long to very long
*Ear - Density: Medium to dense

*Awns or scurs - presence: Present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short

*Awns at tip of ear - length:

Lower glume - Beak length: Short

Lower glume - Beak shape (curvature): Slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow

Lower glume - Shoulder type: Straight or elevated

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak

Lower glume - Internal imprint:

Lower lemma - Beak shape: Slightly curved

Lower lemma - Beak length-: Short

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Weak *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:*Seasonal type:

Winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 31 March 2015

GRAVITAS

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium to dense

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns of scurs - distribution upper ¼

Awns or scurs - length: short

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak lengthmediumBeak curvatureslightly curvedShoulder widthnarrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



HAMPDEN

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs medium

Beak length medium to long
Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late



HARDWICKE

CHARACTERS IN DETAIL

Ear

Colour white
Length medium to long

Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length short to medium
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) weak to medium
Culm - neck weak to medium
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence medium to late



HAVANA

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼
Awns or scurs - length: medium to long

Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak lengthshort to mediumBeak curvaturemoderately curvedShoulder widthnarrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit semi-erect
Time of ear emergence medium



HEREFORD

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium
Density: medium to dense

Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or Scurs - length: long to very long

Apical rachis segment - convex face hairiness: strong

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3

Beak length: medium

Beak shape (curvature): strongly curved

Shoulder medium
Shoulder elevated

Lower Lemma

Beak shape: strongly curved

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: medium Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong

Flag leaf - blade (lower side): strong to very strong

Culm neck: strong
Ear - Glaucosity: strong

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: intermediate Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



HEREWARD

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼

Awns or scurs - length: very short to short

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs weak to medium - Group 1+

Beak lengthmediumBeak curvatureslightly curvedShoulder widthbroad to very broad

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium

Flag leaf - anthocyanin of auricles absent or very weak



HERO

CHARACTERS IN DETAIL

Grain

Colour reddish Colouration with phenol dark

Plant

Anthocyanin colouration of coleoptile weak to medium
Growth habit intermediate
Frequency of plants with recurved flag leaves
Anthocyanin colouration of flag leaf auricles
Time of ear emergence late

Length (stem, ears, awns or scurs) short
Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong

Flag leaf blade (lower side) medium to strong
Ear medium to strong
Culm - neck medium to strong

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium
Shoulder shape slightly elevated
Length of beak medium

Shape of beakmoderately curvedExtent of internal hairsmedium (Group 2)External surface roughnessslightly rough

Ear

Density dense

Length (excluding awns or scurs) short to medium

Awns or scurs - presence scurs present

Awns or scurs - length

Awns or scurs - length long
Colour white
Shape in profile paralle

Shape in profile parallel sided
Apical rachis hairiness on convex surface small to medium
Awns or scurs - distribution upper 1/4



HEROLDO

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: medium
Shape (side profile) tapering

Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short
Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3
Beak length: short to medium

Beak shape (curvature): straight
Shoulder narrow

Shoulder straight to elevated

Lower Lemma

Beak shape:

Grain

Colour white

Colouration with phenol: light to medium

Straw

Length: short to medium

Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong to very strong Flag leaf - blade (lower side): strong to very strong

Culm neck: strong to very strong Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: semi-prostrate Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



HORATIO

CHARACTERS IN DETAIL

Ear

white Colour Length medium lax to medium Density parallel sided Shape-side profile Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/2 Awns or scurs - length: long Apical rachis segment - convex face hairiness medium absent Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length medium to long
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



HUMBER

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼
Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature slightly curved
Shoulder width medium to broad

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence early to medium



Variety Name: **Hybery** crop: Winter Wheat

Breeder's Reference: surh.3628-164 AFP No:

Breeder: Syngenta seeds Ltd Translated from:

Origin: 15 May 2018

Code: HYBE

Coleoptile-anthocyanin colouration: Weak

*Plant - growth habit: Intermediate to semi prostrate

*Time ear emergence: Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Low to medium

*Flag leaf - Glaucosity of sheath: Strong
Flag leaf - Glaucosity of leaf blade: Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Medium to strong

Hairiness of uppermost node:

*Plant - height: Short to medium
*Straw - wall thickness @ 20cms below ear: Very thyin

*Ear - Glaucosity: Medium to strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White
*Ear - Shape (side profile): Parallel sided
Ear - Length: Long

*Ear - Density: Medium to dense *Awns or scurs - presence: Scurs present

Awns or scurs - distibution: *Scurs at tip of ear - length: *Awns at tip of ear - length:

Lower glume - Beak length:Very short to shortLower glume - Beak shape (curvature):Straight to slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width:NarrowLower glume - Shoulder type:Straight

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak to medium

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour:

Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: *Seasonal type:Very dark

Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 16 May 2018



HYPERION

short

CHARACTERS IN DETAIL

Awns or scurs - length

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough
Extent of internal hairs medium (2)
Beak length short to medium
Beak curvature slightly curved
Shoulder width medium

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side)

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence early to medium
Flag leaf - anthocyanin of auricles absent or very weak



Variety Name: **HYTECK** crop: Alternative Wheat

Breeder's Reference: hyteck AFP No:

Breeder: Saaten Union Recherche Translated from:

Origin: Date: 15 May 2018

Code: HYTE

Coleoptile-anthocyanin colouration: Very weak to weak

*Plant - growth habit: Intermediate to semi prostrate

*Time ear emergence: Early to medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Low

*Flag leaf - Glaucosity of sheath: Strong to very strong
Flag leaf - Glaucosity of leaf blade: Medium to strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Medium to strong

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Thin to medium

*Ear - Glaucosity: Medium to strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Parallel sided
Ear - Length: Long to very long

*Ear - Density: Dense

*Awns or scurs - presence: Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short to short

*Awns at tip of ear - length:

Lower glume - Beak length:

Lower glume - Beak shape (curvature):

Very short to short

Straight to slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow

Lower glume - Shoulder type: Sloping or slightly sloping

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Very weak *Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:*Seasonal type:

Alternative

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 16 May 2018



ICEBREAKER

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness strong Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium
Beak curvature slightly curved
Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence



ICON

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns of scurs - distributionupper ¾

Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium to long
Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong

Culm - neck medium

Ear weak to medium

Plant growth characters

Flag leaf - anthocyanin of auricles

Coleoptile - anthocyanin colouration absent or very weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late

(NIAB

INVICTA

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution tip only

Awns or scurs - length: short to medium Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence late



ISTABRAQ

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length short

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect

Time of ear emergence very early to early

Flag leaf - anthocyanin of auricles absent or very weak



JB DIEGO

CHARACTERS IN DETAIL

Ear

Colour white

short to medium Length medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present

Awns of scurs - distribution tip only short to medium Awns or scurs - length:

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length short

Beak curvature slight to moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red Colouration with phenol dark

Straw

short Length Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong weak to medium

Ear

Plant growth characters

Coleoptile - anthocyanin colouration strong

semi-erect to intermediate Plant - growth habit

Time of ear emergence medium



JELLICOE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present

Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth Extent of internal hairs medium (2)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



JORVIK

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium
Beak length medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape straight

Lower lemma

Beak curvature geniculate

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late



JOYFUL

CHARACTERS IN DETAIL

Ear

Colour white
Length long
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



variety Name: Juniper crop: Wheat

Breeder's Reference: SjP0579 **AFP No:**

Breeder: Limagrain UK Ltd Translated from: Danish

Origin: Date: 06 April 2023

Code: JUNP

Coleoptile-anthocyanin colouration: Medium

*Plant - growth habit: Intermediate to semi-prostrate

*Time ear emergence: Medium

Flag leaf - Anthocyanin colouration of auricles:

Absent or very weak
Frequency of plants with recurved flag leaves:

Low to medium

*Flag leaf - Glaucosity of sheath: Strong
Flag leaf - Glaucosity of leaf blade: Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Medium to strong

Hairiness of uppermost node:

*Plant - height: Medium

*Straw - wall thickness @ 20cms below ear: Thin

*Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White
*Ear - Shape (side profile): Parallel sided

Ear - Length: Short to medium

*Ear - Density: Medium

*Awns or scurs - presence: Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: hort to medium

*Awns at tip of ear - length:

Lower glume - Beak length: Long

Lower glume - Beak shape (curvature):Slightly curved to moderately curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow to medium

Lower glume - Shoulder type:Horizontal to slightly elevated

Lower glume - External surface roughness: Lower glume - Extent of internal hairs: Lower glume - Internal imprint:

Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour: Reddish

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type: Winter type

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 19 April 2023



KETCHUM

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



KINGDOM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution tip only
Awns or scurs - length: short
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



KINGFISHER

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium to long
Beak curvature slightly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KIPLING

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitydense to very denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper ½

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak lengthmediumBeak curvatureslightly curvedShoulder widthmediumShoulder shapeelevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side)

Culm - neck medium Ear weak

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



KRIS

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium

Density: medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short
Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3

Beak length: short to medium

Beak shape (curvature): slightly curved

Shoulder medium

Shoulder straight to elevated

Lower Lemma

Beak shape: slightly curved

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: short Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong to very strong Flag leaf - blade (lower side): medium to strong

Culm neck: strong Ear - Glaucosity: strong

Plant growth characters

Coleoptile-anthocyanin medium to strong

Plant - growth habit: semi-prostrate to prostrate

Time ear emergence: late

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



KWS ALABASTER

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present

Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium (group 2)

Beak length short

Beak curvature slightly curved Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour white

Colouration with phenol medium to dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence early to medium



KWS ARNIE

CHARACTERS IN DETAIL

Grain

Colour reddish Colouration with phenol medium

Plant

Anthocyanin colouration of coleoptile medium to strong
Growth habit intermediate

Frequency of plants with recurved flag leaves
Anthocyanin colouration of flag leaf auricles
Time of ear emergence
Length (stem, ears, awns or scurs)

low
absent or very weak
medium to late
very short to short

Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong
Flag leaf blade (lower side) strong to very strong
Ear strong to very strong
Culm - neck strong to very strong

Lower glume

Hairiness of external surface absent Shoulder width narrow

Shoulder shape horizontal to slightly elevated

Length of beak medium

Shape of beak slight to moderately curved

Extent of internal hairs medium (Group 2)

External surface roughness rough

Ear

Density medium to dense
Length (excluding awns or scurs) medium to long
Awns or scurs - presence scurs present
Awns or scurs - length long

Colour white
Shape in profile tapering

Apical rachis hairiness on convex surface small to medium Awns or scurs - distribution upper 3/4



KWS BARREL

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length medium

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence late to very late



KWS BASSET

medium to strong

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length long

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) weak to medium

Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate
Time of ear emergence late to very late



KWS BIELSUM

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitydenseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



KWS BOATUM

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long

Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long
Beak curvature strongly curved
Shoulder width narrow to medium
Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-prostrate

Time of ear emergence late to very late

Flag leaf - anthocyanin of auricles absent or very weak



KWS BONHAM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper ¼

Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) weak to medium

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS BRENNUM

CHARACTERS IN DETAIL

Ear

Colour white Length long

Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs weak to medium (group 1+)

Beak length short

Beak curvature slight to moderately curved

Shoulder width medium

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol light to medium

Straw

Length short

Wall thickness @20cms below ear

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS BRIUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long

Apical rachis segment - convex face hairiness strong to very strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS BURGHLEY

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium
Shape-side profile tapering

Awns or scurs - presence scurs present

Awns or scurs - distribution upper 1/4

Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS CASHEL

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense

Shape-side profile tapering

Awns or scurs - presence awns present
Awns or scurs - distribution whole length
Awns or scurs - length very long
Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth Extent of internal hairs strong

Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath weak

Flag leaf- blade (lower side) very weak to weak

Culm - neck weak Ear weak

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late



KWS CHILOE

CHARACTERS IN DETAIL

Ear

white Colour Length medium lax to medium Density Shape-side profile tapering Awns or scurs - presence awns present Awns of scurs - distribution whole length Awns or scurs - length: very long Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairsstrong - Group 3Beak lengthlong to very longBeak curvatureslightly curved

Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol very light to light

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) medium to strong

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



KWS CLEVELAND

long

CHARACTERS IN DETAIL

Awns or scurs - length:

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper ¼

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit semi-erect
Time of ear emergence medium



Variety Name: KWS COLOSSEUM crop: Wheat

Breeder's KWS W370 AFP No:

Breeder: KWS UK LTD Translated from:

Origin: Date: 28 June 2021

Code: KCOL

Coleoptile-anthocyanin colouration: Medium to strong

*Plant - growth habit: Intermediate to semi-prostrate

*Time ear emergence: Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Low

*Flag leaf - Glaucosity of sheath: Strong to very strong
Flag leaf - Glaucosity of leaf blade: Strong to very strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong to very strong

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Absent or very thin

*Ear - Glaucosity: Strong to very strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White
*Ear - Shape (side profile): Parallel sided

Ear - Length: Medium

*Ear - Density: Medium to dense *Awns or scurs - presence: Scur present

Awns or scurs - distibution: *Scurs at tip of ear - length:

*Awns at tip of ear - length: Medium to long

Lower glume - Beak length: Medium

Lower glume - Beak shape (curvature):Slightly curved to moderately curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Medium to broad

Lower glume - Shoulder type: Slightly sloping to straight

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Medium

Lower glume - Internal imprint: Slightly curved to moderately curved

Lower lemma - Beak shape: Strongly curved to geniculate

Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Medium
*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: Dark to very dark

*Seasonal type: Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 28 June 2021



KWS CONROS

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium (group 1+)

Beak lengthshort to mediumBeak curvatureslightly curvedShoulder widthnarrowShoulder shapeelevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong

Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence very late



KWS COYNE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length long
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium to long
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak Plant - growth habit intermediate

late

Time of ear emergence



KWS CRANIUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature slightly curved

Shoulder width narrow

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS CRISPIN

medium to strong

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthshort to medium

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs medium (2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



KWS CROFT

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼
Awns or scurs - length: long
Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



KWS CROSBY

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper ½ Awns or scurs - length: long weak

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2 medium to long Beak length Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong

Culm - neck strong to very strong Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



KWS DALI

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns of scurs - distributionupper ½Awns or scurs - length:medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf - blade (lower side) weak to medium
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS DAWSUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



KWS DELUXUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium (group 2)
Beak length short to medium
Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



KWS DINKUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence lat

Flag leaf - anthocyanin of auricles absent or very weak



KWS DRAGUM

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense
Shape-side profile parallel sided

Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish
Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) medium to strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS EQUIPE

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol medium to dark

Plant

Anthocyanin colouration of coleoptile weak

Growth habit semi-erect to intermediate

Frequency of plants with recurved flag leaves
Anthocyanin colouration of flag leaf auricles
Time of ear emergence
low to medium
absent or very weak
medium to late

Length (stem, ears, awns or scurs) short Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong

Flag leaf blade (lower side) strong
Ear strong
Culm - neck strong

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium
Shoulder shape slightly elevated
Length of beak medium

Shape of beak moderately curved
Extent of internal hairs medium (Group 2)
External surface roughness smooth

Ear

Density medium
Length (excluding awns or scurs) medium
Awns or scurs - presence scurs present

Awns or scurs - length long
Colour white
Shape in profile tapering

Apical rachis hairiness on convex surface very small to small

Awns or scurs - distribution upper 3/4

WINTER WHEAT



KWS EVOKE

CHARACTERS IN DETAIL

Ear

Colour white

Length medium

Density medium

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns of scurs - distribution upper ¼

Awns or scurs - length: short to medium
Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length short

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium

Flag leaf - anthocyanin of auricles

WINTER WHEAT



KWS EVOLVE

CHARACTERS IN DETAIL

Ear

white Colour Length long Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 medium Awns or scurs - length Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs weak to medium
Beak length medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colourationabsent to weakPlant - growth habitintermediateTime of ear emergencelate



KWS EXTASE

CHARACTERS IN DETAIL

Ear

Colour white
Length medium to long

Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length short

Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate
Time of ear emergence medium



KWS FENWAY

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature INVALID CHARACTER STATE!

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate

Time of ear emergence very late Flag leaf - anthocyanin of auricles



KWS FIREFLY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to longApical rachis segment - convex face hairinessmedium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate
Time of ear emergence late



KWS FLUTE

CHARACTERS IN DETAIL

Grain

Colour reddish
Colouration with phenol dark

Plant

Anthocyanin colouration of coleoptile weak to medium intermediate

Frequency of plants with recurved flag leaves absent or very low absent or very weak

Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late

Length (stem, ears, awns or scurs) very short to short

Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong
Flag leaf blade (lower side) strong to very strong
Ear strong to very strong
Culm - neck strong to very strong

Lower glume

Hairiness of external surface absent Shoulder width narrow

Shoulder shape horizontal to slightly elevated

Length of beak medium

Shape of beak slight to moderately curved

Extent of internal hairs very large (Group 3)

External surface roughness smooth

Ear

Density dense
Length (excluding awns or scurs) medium
Awns or scurs - presence scurs present
Awns or scurs - length long
Colour white

Shape in profile parallel sided
Apical rachis hairiness on convex surface medium to large
Awns or scurs - distribution upper 1/2



KWS FOXTUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs weak (group 1)
Beak length medium

Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape slightly elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



KWS FULCRUM

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong

Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS GATOR

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½
Awns or scurs - length: long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence medium to late



KWS GIGANTUM

absent

CHARACTERS IN DETAIL

Anthers - anthocyanin colouration

Ear

white Colour medium to long Length Density medium to dense Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness large

Lower glume

External surface rough

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish
Colouration with phenol medium to dark

colouration man promo.

Straw
Length short
Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS GUIUM

CHARACTERS IN DETAIL

Ear

Colour white Length long Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 3/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

medium to strong (group 2+) Extent of internal hairs

Beak length short to medium Beak curvature strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) medium to strong Culm - neck strong to very strong medium

Ear

Plant growth characters

Coleoptile - anthocyanin colouration strong Plant - growth habit intermediate Time of ear emergence late to very late Flag leaf - anthocyanin of auricles absent or very weak



KWS HARLYN

medium

CHARACTERS IN DETAIL

Awns or scurs - length

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface rough
Extent of internal hairs medium (2)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



KWS HENUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long

Apical rachis segment - convex face hairiness strong to very strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length long

Beak curvature strongly curved
Shoulder width medium to broad
Shoulder shape elev. to strongly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermediate

Time of ear emergence late to very late

Flag leaf - anthocyanin of auricles absent or very weak



KWS HORIZON

CHARACTERS IN DETAIL

Ear

Colour white

Length medium

Density medium

Shape-side profile tapering

Awns or scurs - presence scurs present

Awns of scurs - distribution upper ½

Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak lengthmediumBeak curvatureslightly curvedShoulder widthnarrowShoulder shapeelevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



KWS JACKAL

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence medium to late



KWS JUBILATION

CHARACTERS IN DETAIL

Ear

white Colour Length medium lax to medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length short to medium
Beak curvature strongly curved
Shoulder width narrow

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS JUGUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length lon

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS KARTUM

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium to dense

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns or scurs - distribution upper 1/2

Awns or scurs - length long to very long

Apical rachis segment - convex face hairiness large
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length short to medium
Beak curvature moderately curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish
Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS KENTON

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns of scurs - distribution upper ½ Awns or scurs - length: medium to long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) weak to medium
Culm - neck weak to medium

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS KERANKUM

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) strong
Culm - neck very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



KWS KERRIN

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present upper 1/4 Awns or scurs - distribution Awns or scurs - length medium to long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) weak to medium
Culm - neck weak to medium
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate Time of ear emergence very late



KWS KIELDER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns of scurs - distributionupper ½

Awns or scurs - length: long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium to long
Beak curvature slightly curved

Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit semi-prostrate Time of ear emergence medium to late



KWS KINETIC

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long
Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS KINNELLUM

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermediate

Time of ear emergence medium to late

Flag leaf - anthocyanin of auricles absent or very weak



KWS LEON

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs weak to medium (1+)
Beak length short to medium
Beak curvature moderately curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence late to very late



KWS LILI

CHARACTERS IN DETAIL

Ear

white Colour

Length short to medium

Density dense Shape-side profile

parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4

Awns or scurs - length medium Apical rachis segment - convex face hairiness weak absent

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs medium

Beak length short to medium Beak curvature slightly curved Shoulder width narrow

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

short Length

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



KWS LUMOS

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼
Awns or scurs - length: medium
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium
Beak curvature strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colourationstrongPlant - growth habitsemi-erectTime of ear emergencemedium



KWS LUTHER

CHARACTERS IN DETAIL

Ear

white Colour Length medium lax to medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 3/4 Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium



KWS MADRYN

CHARACTERS IN DETAIL

Ear

Colour white

Length medium

Density medium

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns of scurs - distribution upper ½

Awns or scurs - length: medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



KWS MCGUGUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness strong to very strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length short to medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



KWS MODELLUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length very long
Apical rachis segment - convex face hairiness large
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-prostrate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



KWS MOJITO

weak to medium

late

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4Awns or scurs - lengthlong to very long

Anthers - anthocyanin colouration absent

Apical rachis segment - convex face hairiness

Lower glume

External surface smooth

Extent of internal hairs weak to medium

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) weak to medium

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate

Time of ear emergence



KWS MONGOOSE

CHARACTERS IN DETAIL

Grain

Colour reddish Colouration with phenol dark

Plant

Anthocyanin colouration of coleoptile weak to medium Growth habit intermediate Frequency of plants with recurved flag leaves very low to low Anthocyanin colouration of flag leaf auricles absent or very weak Time of ear emergence medium Length (stem, ears, awns or scurs) very short to short

Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong Flag leaf blade (lower side) strong to very strong strong to very strong Culm - neck strong to very strong

Lower glume

Hairiness of external surface absent Shoulder width narrow Shoulder shape slightly elevated Length of beak medium

Shape of beak moderate to strongly curved Extent of internal hairs very large (Group 3)

External surface roughness smooth

Ear

Density dense Length (excluding awns or scurs) medium Awns or scurs - presence scurs present Awns or scurs - length medium to long Colour white

Shape in profile parallel sided Apical rachis hairiness on convex surface small to medium

Awns or scurs - distribution upper 1/2



Variety Name: KWS Montana crop: Winter wheat

Breeder's Reference: KW 8027-3-08 AFP No:

Breeder: KWS LOCHOW GMBH Translated from: German

Origin: Date: 22 March 2016

Code: KWMON

Coleoptile-anthocyanin colouration:Medium*Plant - growth habit:Semi prostrate

*Time ear emergence: Late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Absent or very low

*Flag leaf - Glaucosity of sheath: Strong
Flag leaf - Glaucosity of leaf blade: Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong to very strong

Hairiness of uppermost node:

*Plant - height: Short to medium
*Straw - wall thickness @ 20cms below ear: Absent or very thin

*Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile):

Ear - Length:

*Ear - Density:

*Awns or scurs - presence:

Parallel sided

Medium

Medium

Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short

*Awns at tip of ear - length:

Lower glume - Beak length:

Lower glume - Beak shape (curvature):

Very short to short

Straight to slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Medium

Lower glume - Shoulder type: Slightly sloping to straight

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Very weak to weak

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Strong to very strong

*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:Medium to dark

*Seasonal type: Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 23 March 2016



KWS MOZART

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium
Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



KWS NEWBIE

CHARACTERS IN DETAIL

Grain

Colour reddish Colouration with phenol medium

Plant

Anthocyanin colouration of coleoptile weak to medium
Growth habit intermediate
Frequency of plants with recurved flag leaves
Anthocyanin colouration of flag leaf auricles absent or very weak
Time of ear emergence medium to late

Length (stem, ears, awns or scurs) short Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong

Flag leaf blade (lower side) strong
Ear strong
Culm - neck strong

Lower glume

Hairiness of external surface absent Shoulder width narrow

Shoulder shape slightly elevated to strongly elevated

Length of beakmediumShape of beakstrongly curvedExtent of internal hairsmedium (Group 2)

External surface roughness entire external surface rough

Ear

Density medium to dense
Length (excluding awns or scurs) short to medium
Awns or scurs - presence scurs present
Awns or scurs - length medium
Colour white

Shape in profile tapering
Awns or scurs - distribution upper 1/4



KWS PALLADIUM

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns or scurs - distribution upper 1/2

Awns or scurs - length long

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature moderately curved
Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS PARKIN

CHARACTERS IN DETAIL

Ear

Colour white
Length short
Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration

Lower glume

External surface entire external surface rough Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature moderately curved Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol light to medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



KWS PLECTRUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitydenseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length short to medium
Beak curvature v.slightly curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong
Ear very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS PODIUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit semi-erect
Time of ear emergence medium



KWS RACKUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long
Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS ROADUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long

Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape horizontal to slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS ROWAN

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ½

Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium



KWS SABRUM

CHARACTERS IN DETAIL

Ear

Colour white
Length short
Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs strong (group 3)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong

Flag leaf- blade (lower side) strong to very strong

Culm - neck very strong Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium

Flag leaf - anthocyanin of auricles absent or very weak



KWS SANTIAGO

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼
Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



KWS SAXTEAD

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence awns present
Awns of scurs - distribution whole length
Awns or scurs - length:

Awns or scurs - length: long
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs strong - Group 3

Beak length long

Beak curvature strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit semi-erect
Time of ear emergence medium



KWS SCIMITAR

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width broad

Shoulder shape straight to elevated

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate
Time of ear emergence late



KWS SCOPE

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol light to medium

Plant

Anthocyanin colouration of coleoptile strong

intermed. to semi-prostrate Growth habit

Frequency of plants with recurved flag leaves very low to low Anthocyanin colouration of flag leaf auricles absent or very weak Time of ear emergence medium to late

Length (stem, ears, awns or scurs) short

Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong Flag leaf blade (lower side) strong to very strong

strong to very strong Culm - neck strong to very strong

Lower glume

Hairiness of external surface absent

Shoulder width very narrow to narrow Shoulder shape slightly elevated Length of beak short to medium Shape of beak moderately curved very large (Group 3)

Extent of internal hairs

External surface roughness smooth

Ear

Density dense

Length (excluding awns or scurs) medium Awns or scurs - presence scurs present Awns or scurs - length medium white

Colour

Shape in profile parallel sided Apical rachis hairiness on convex surface very small to small

Awns or scurs - distribution upper 1/4

KWS SILVERSTONE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs medium (2)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



KWS SILVUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong

Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



KWS SISKIN

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium (1+)
Beak length short to medium

Beak curvature slight to moderately curved Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



KWS SKATEUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-prostrate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



KWS SKOMER

medium to long

CHARACTERS IN DETAIL

Awns or scurs - length

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak lengthmedium to longBeak curvaturestrongly curvedShoulder widthnarrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol light to medium

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence early



KWS SOLITAIRE

CHARACTERS IN DETAIL

Grain

Colour reddish Colouration with phenol dark

Plant

Anthocyanin colouration of coleoptile very weak to weak Growth habit intermediate
Frequency of plants with recurved flag leaves very low to low Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late
Length (stem, ears, awns or scurs) short
Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong
Flag leaf blade (lower side) strong to very strong
Ear strong to very strong
Culm - neck strong to very strong

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium

Shoulder shape horizontal to slightly elevated

Length of beak medium

Shape of beak moderate to strongly curved

Extent of internal hairs very large (Group 3)

External surface roughness smooth

Ear

Density dense

Length (excluding awns or scurs)short to mediumAwns or scurs - presencescurs presentAwns or scurs - lengthvery longColourwhite

Shape in profile parallel sided
Apical rachis hairiness on convex surface medium
Awns or scurs - distribution upper 3/4



KWS SOLO

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns of scurs - distributionupper ¼Awns or scurs - length:medium to long

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong

Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate
Time of ear emergence medium to late



KWS STERLING

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns of scurs - distribution upper ¼
Awns or scurs - length: long
Apical rachis segment - convex face hairiness weak

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved
Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



KWS TARGET

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense
Shape-side profile parallel sided

Awns or scurs - presence scurs present
Awns or scurs - distribution upper ¼
Awns or scurs - length: medium
Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit semi-erect
Time of ear emergence medium



KWS TEMPO

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium
Beak length short to medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence very late



KWS TRAINUM

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length long

Beak curvature strongly curved to geniculate

Shoulder width medium

Shoulder shape slightly elevated

Lower lemma

Beak curvature geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-prostrate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



KWS TRINITY

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs weak to medium
Beak length medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence late



KWS TROMMEL

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong to very longApical rachis segment - convex face hairinessmedium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature strongly curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence very late



KWS ULTIMATUM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs strong (group 3)
Pack langth

Beak lengthmediumBeak curvaturestrongly curvedShoulder widthvery narrow to narrowShoulder shapeslightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS VICARAGE

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol dark to very dark

Plant

Anthocyanin colouration of coleoptile weak
Growth habit intermediate

Frequency of plants with recurved flag leaves low

Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late

Length (stem, ears, awns or scurs) very short to short

Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong
Flag leaf blade (lower side) strong

Ear medium to strong

Culm - neck strong

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium

Shoulder shape horizontal to slightly elevated

Length of beak medium to long

Shape of beak moderate to strongly curved

Extent of internal hairs very large (Group 3)

External surface roughness smooth

Ear

Density medium to dense
Length (excluding awns or scurs) short to medium
Awns or scurs - presence scurs present
Awns or scurs - length medium to long

Colour white

Shape in profile parallel sided
Apical rachis hairiness on convex surface small to medium

Awns or scurs - distribution upper 1/2



KWS VINNIE

CHARACTERS IN DETAIL

Grain

Colour reddish
Colouration with phenol dark

Plant

Anthocyanin colouration of coleoptile medium

Growth habit intermed. to semi-prostrate

Frequency of plants with recurved flag leaves very low to low Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late
Length (stem, ears, awns or scurs) short
Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong
Flag leaf blade (lower side) strong
Ear strong
Culm - neck strong

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium
Shoulder shape slightly elevated
Length of beak short to medium

Shape of beak slight to moderately curved Extent of internal hairs very large (Group 3)

External surface roughness smooth

Ear

Density dense
Length (excluding awns or scurs) medium
Awns or scurs - presence scurs present
Awns or scurs - length short to medium

Colour white

Shape in profile parallel sided
Apical rachis hairiness on convex surface medium
Awns or scurs - distribution upper 1/2



KWS WEBBUM

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long to very long Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak lengthmedium to longBeak curvaturestrongly curvedShoulder widthnarrow to mediumShoulder shapeslightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



KWS WRENUM

CHARACTERS IN DETAIL

Ear

white Colour medium Length Density dense Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness large to very large

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium to long
Beak curvature moderately curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish
Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



KWS YARIS

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium to long

Density: medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: medium

Apical rachis segment - convex face hairiness: medium to strong

Anthers - anthocyanin colouration

Lower Glume

External surface roughness:

Extent of internal hairs: strong - group 3

Beak length: long

Beak shape (curvature): strongly curved

Shoulder medium
Shoulder elevated

Lower Lemma

Beak shape: strongly curved

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: medium Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong to very strong

Flag leaf - blade (lower side): very strong

Culm neck: strong to very strong
Ear - Glaucosity: strong to very strong

Plant growth characters

Coleoptile-anthocyanin absent or very weak Plant - growth habit: semi-prostrate

Time ear emergence: late

Flag leaf - Anthocyanin colouration of auricles:



KWS ZENTARR

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long to very long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence late to very late



KWS ZYATT

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs strong (group 3)

Beak length short to medium

Beak curvature slightly curved

Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence very late



LANCASTER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ¼Awns or scurs - length:long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck weak to medium

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence



LANGDALE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough
Extent of internal hairs medium (2)
Beak length short to medium
Beak curvature slightly curved
Shoulder width medium

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit semi-erect
Time of ear emergence late



LEAR

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ¼
Awns or scurs - length: medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



LEEDS

long

CHARACTERS IN DETAIL

Awns or scurs - length:

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper ½

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium



LG ALDANITI

CHARACTERS IN DETAIL

Ear

Colour white Length short

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence awns present
Awns or scurs - distribution whole length
Awns or scurs - length very long

Apical rachis segment - convex face hairiness very small to small

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length long to very long

Beak curvature moderate to strongly curved Shoulder width very narrow to narrow

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature

Grain

Colour reddish
Colouration with phenol very dark

Straw

Length very short to short Wall thickness @20cms below ear thick or filled

Glaucosity

Flag leaf - sheath absent or very weak Flag leaf- blade (lower side) absent or very weak Culm - neck absent or very weak Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles very weak



LG ARKLE

absent

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length long
Apical rachis segment - convex face hairiness large

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs strong (group 3)

Beak length medium to long
Beak curvature v.slightly curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence lat



LG ASTRONOMER

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density lax to medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long to very long

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late



LG BEOWULF

long

CHARACTERS IN DETAIL

Ear

white Colour

Length medium to long Density medium to dense Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs strong (group 3) Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak Plant - growth habit intermediate

Time of ear emergence late



LG BLETCHLEY

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns or scurs - distribution upper 1/4

Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



LG CASSIDY

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath absent or very weak Flag leaf- blade (lower side) absent or very weak Culm - neck absent or very weak Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



LG DETROIT

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late



LG FARRIER

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4

Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface entire external surface rough
Extent of internal hairs medium to strong (group 2+)

Extent of internal hairs medium to strong (group 2+)
Beak length medium

Beak curvature strongly curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) strong
Culm - neck very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate

Time of ear emergence late



LG GENERATION

long

CHARACTERS IN DETAIL

Awns or scurs - length

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) weak

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence medium to late



LG GILMORE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitydenseShape-side profileparallel sided

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution tip only

Awns or scurs - length very short to short
Apical rachis segment - convex face hairiness large to very large

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak lengthvery short to shortBeak curvatureslightly curvedShoulder widthnarrow to mediumShoulder shapeslightly elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour reddish

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate



LG GRADUATE

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length lon

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late to very late



LG GRENDEL

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)

Beak length medium to long

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late



LG HENRI

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol dark to very dark

Plant

Anthocyanin colouration of coleoptile medium to strong
Growth habit intermediate
Frequency of plants with recurved flag leaves absent or very low
Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late
Length (stem, ears, awns or scurs) short
Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong
Flag leaf blade (lower side) strong to very strong
Ear strong to very strong
Culm - neck strong to very strong

Lower glume

Hairiness of external surface absent
Shoulder width narrow
Shoulder shape slightly elevated

Length of beak long

Shape of beak moderate to strongly curved
Extent of internal hairs very small to medium (Group 1+)

External surface roughness slightly rough

Ear

Density medium to dense
Length (excluding awns or scurs) medium

Awns or scurs - presence scurs present

Awns or scurs - length long to very long

Colour white

Shape in profile parallel sided
Apical rachis hairiness on convex surface medium
Awns or scurs - distribution upper 3/4



LG ILLUMINATE

CHARACTERS IN DETAIL

Ear

Colour white
Length medium to long
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4

Awns or scurs - distribution upper 1/4
Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs medium (group 2)

Beak length long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late



LG INTERSTELLAR

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs medium (group 2)
Beak length medium to long
Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence late



LG JIGSAW

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 3/4 Awns or scurs - length long Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slightly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence late to very late



LG KERMIT

CHARACTERS IN DETAIL

Grain

Colour reddish Colouration with phenol dark

Plant

Anthocyanin colouration of coleoptile very weak to weak Growth habit intermediate

Frequency of plants with recurved flag leaves low
Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence lat

Length (stem, ears, awns or scurs) very short to short

Glaucosity

Flag leaf sheath medium to strong
Flag leaf blade (lower side) medium to strong
Ear weak to medium
Culm - neck medium to strong

Lower glume

Hairiness of external surface absent

Shoulder width very narrow to narrow Shoulder shape slightly elevated Length of beak short to medium Shape of beak slightly curved Extent of internal hairs very small (Group 1)

External surface roughness smooth

Ear

Density medium

Length (excluding awns or scurs) medium to long

Awns or scurs - presence scurs present

Awns or scurs - length short

Colour white

Shape in profile tapering
Awns or scurs - distribution upper 1/4



LG MEDUSA

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length long

Beak curvature strongly curved
Shoulder width very narrow to narrow
Shoulder shape elev. to strongly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



LG MOTOWN

CHARACTERS IN DETAIL

Ear

white Colour Length short Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs strong (group 3)
Beak length short to medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence late



LG NASHVILLE

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length short to medium medium

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs strong (group 3) Beak length medium Beak curvature slightly curved Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

very short to short Length Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath absent or very weak Flag leaf- blade (lower side) absent or very weak Culm - neck absent or very weak Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate late

Time of ear emergence



LG PARTRIDGE

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol medium to dark

Plant

Anthocyanin colouration of coleoptile medium to strong

Growth habit intermed. to semi-prostrate

Frequency of plants with recurved flag leaves very low to low Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late Length (stem, ears, awns or scurs) short Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong

Flag leaf blade (lower side) strong Ear strong Culm - neck strong

Lower glume

Hairiness of external surface absent Shoulder width narrow

Shoulder shape slightly elevated Length of beak short to medium

Shape of beak slight to moderately curved Extent of internal hairs very large (Group 3)

smooth

External surface roughness

Ear

medium to dense Density

medium Length (excluding awns or scurs) Awns or scurs - presence scurs present Awns or scurs - length medium to long

Colour white

Shape in profile parallel sided Apical rachis hairiness on convex surface small to medium Awns or scurs - distribution upper 1/4



LG PHAROS

CHARACTERS IN DETAIL

Ear

Colour white Length long

Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long
Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak Plant - growth habit intermediate

Time of ear emergence late



LG PRINCE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong to very long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak (group 1)

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



LG PUFFIN

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol light to medium

Plant

Anthocyanin colouration of coleoptile medium

Growth habit intermediate

Frequency of plants with recurved flag leaves low
Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late

Length (stem, ears, awns or scurs) very short to short

Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong

Flag leaf blade (lower side) strong
Ear strong
Culm - neck strong

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium

Shoulder shape horizontal to slightly elevated

Length of beak short to medium

Shape of beak moderate to strongly curved
Extent of internal hairs very small to medium (Group 1+)

External surface roughness smooth

Ear

Density dense

Length (excluding awns or scurs)short to mediumAwns or scurs - presencescurs presentAwns or scurs - lengthmedium to long

Colour white

Shape in profile parallel sided
Apical rachis hairiness on convex surface small to medium
Awns or scurs - distribution upper 1/4



WINTER WHEAT

AFP 1/2836

LG QUASAR

CHARACTERS IN DETAIL

Ear

Colour white Length medium Density dense parallel sided Shape-side profile Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width narrow to medium Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late



Winter Wheat

AFP 1/3247

LG REBELLION

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol medium to dark

Plant

Anthocyanin colouration of coleoptile

Growth habit

Frequency of plants with recurved flag leaves

Anthocyanin colouration of flag leaf auricles

Time of ear emergence

Length (stem, ears, awns or scurs)

Straw wall thickness (20cm below ear)

medium to strong intermediate

very low to low

absent or very weak medium

very short to short

thin

Glaucosity

Flag leaf sheath

Flag leaf blade (lower side)

Ear

Culm - neck

strong to very strong $% \left(t\right) =\left(t\right) \left(t\right) \left($

strong to very strong

strong to very strong strong to very strong

Lower glume

Hairiness of external surface

Shoulder width

Shoulder shape

Length of beak

Shape of beak

Extent of internal hairs

External surface roughness

absent

narrow to medium

horizontal to slightly elevated

short

v.slightly curved

medium (Group 2)

smooth

Ear

Density

Length (excluding awns or scurs)

Awns or scurs - presence Awns or scurs - length

Colour

Shape in profile

Apical rachis hairiness on convex surface

Awns or scurs - distribution

medium medium

scurs present

medium white

tapering

medium to large

upper 1/2



LG REDRUM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4Awns or scurs - lengthlong

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length mediun

Beak curvature moderate to strongly curved

Shoulder width medium

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish
Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late
Flag leaf - anthocyanin of auricles very weak



LG REDWALD

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long to very long

Apical rachis segment - convex face hairiness large
Anthers - anthocyanin colouration absent

Lower glume

External surface

Extent of internal hairs strong (group 3)
Beak length medium to long
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol dark to very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate

Time of ear emergence late



LG RHYTHM

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs weak (group 1)
Beak length medium

Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late



LG ROCKHOPPER

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium
Shape-side profile tapering
Awns or scurs - presence awns present

Awns or scurs - distribution whole length
Awns or scurs - length very long

Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs strong (group 3)
Beak length very long

Beak curvature moderate to strongly curved

Shoulder width narrow

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) medium to strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles very weak



LG SABERTOOTH

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution

Awns or scurs - length long
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium to long
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



LG SEEKER

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4

Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough Extent of internal hairs medium to strong (group 2+)

Beak length medium
Beak curvature strongly curved
Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



LG SHERGAR

very low to low

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol medium to dark

Plant

Anthocyanin colouration of coleoptile very weak to weak intermediate

Growth habit

Frequency of plants with recurved flag leaves

Time of ear emergence

Length (stem, ears, awns or scurs)

Straw wall thickness (20cm below ear)

Anthocyanin colouration of flag leaf auricles absent or very weak medium to late very short thin

Glaucosity

Flag leaf sheath strong to very strong

Flag leaf blade (lower side) strong to very strong strong to very strong

Culm - neck strong to very strong

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium

Shoulder shape slightly elevated to strongly elevated

Length of beak medium Shape of beak strongly curved

Extent of internal hairs medium (Group 2)

External surface roughness smooth

Ear

Density medium to dense Length (excluding awns or scurs) medium to long

Awns or scurs - presence scurs present Awns or scurs - length medium to long

Colour white Shape in profile tapering

Apical rachis hairiness on convex surface small to medium

Awns or scurs - distribution upper 1/2



LG SKYSCRAPER

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long to very long weak to medium

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

red

medium to dark Colouration with phenol

Straw

short to medium Length Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence medium to late



LG SPOTLIGHT

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs weak (group 1)
Beak length medium

Beak length mediur
Beak curvature slight t

Beak curvature slight to moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

GlaucosityFlag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate

Time of ear emergence late



LG SUNDANCE

CHARACTERS IN DETAIL

Ear

white Colour Length short Density dense Shape-side profile parallel sided scurs present Awns or scurs - presence Awns or scurs - distribution upper 1/4 Awns or scurs - length long to very long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs strong (group 3)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) weak
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence very late



LG TAPESTRY

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong

Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



LG TICKET

CHARACTERS IN DETAIL

Ear

Colour white Length short

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs whole internal surface covered

Beak length medium to long
Beak curvature strongly curved
Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



LG TYPHOON

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong

Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



LG VOODOO

medium

CHARACTERS IN DETAIL

Awns or scurs - length

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath absent or very weak
Flag leaf- blade (lower side) absent or very weak
Culm - neck absent or very weak
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium to late

Flag leaf - anthocyanin of auricles absent or very weak



Lion

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4

Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium to strong (group 2+)

Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape horizontal to slightly elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour reddish

Colouration with phenol dark to very dark

Straw

Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence early to medium



LOXTON

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) medium to strong
Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence medium to late

Flag leaf - anthocyanin of auricles absent or very weak



LUSTRE

CHARACTERS IN DETAIL

Ear

white Colour Length medium lax to medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)

Beak length short

Beak curvature slightly curved Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour white

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermediate
Time of ear emergence medium



MAGISTER

CHARACTERS IN DETAIL

Ear

Colour: white
Length: medium
Density: lax to medium
Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: weak to medium -group 1 to 2

Beak length: short

Beak shape (curvature): moderately curved
Shoulder medium to broad

Shoulder straight

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: dark

Straw

Length: medium to long Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): strong

Culm neck: strong

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin weak

Plant - growth habit: semi-prostrate
Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



MARIBOSS

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthshortDensitymediumShape-side profiletaperingAwns or scurs - presencescurs present

Awns or scurs - distribution

Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface
Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) strong to very strong

Culm - neck strong

Ear very weak to weak

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence early



MARIS HUNTSMAN

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs weak to medium (group 1+)

Beak lengthmedium to longBeak curvatureslightly curvedShoulder widthmediumShoulder shapeelevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length long to very long Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium
Flag leaf - anthocyanin of auricles weak



MARIS WIDGEON

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensityvery lax to laxShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper ¼

Awns or scurs - length: medium to long

Anical rachis segment - convex face hairiness medium

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak - Group 1

Beak length medium

Beak curvature v.slightly curved

Beak curvature v.slightly curved
Shoulder width narrow to medium

Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very long Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate

Time of ear emergence late Flag leaf - anthocyanin of auricles weak



Variety Name: Marlowe

Crop:

Breeder's Reference: Sj9726005 AFP No: 1/2413

Breeder: Senova Ltd Translated from: Danish

Origin: Date: 23 March 2017

Code: MARL

Coleoptile-anthocyanin colouration:Weak to medium*Plant - growth habit:Semi prostrate

*Time ear emergence: Late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Medium to high

*Flag leaf - Glaucosity of sheath: Strong
Flag leaf - Glaucosity of leaf blade: Strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong

Hairiness of uppermost node:

*Plant - height: Medium

*Straw - wall thickness @ 20cms below ear: Absent or very thin

*Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White *Ear - Shape (side profile): Parallel

Ear - Length:Short to medium*Ear - Density:Dense to very dense

*Awns or scurs - presence: Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Medium
*Awns at tip of ear - length: Medium
Lower glume - Beak length: Long

Lower glume - Beak shape (curvature): Strongly curved

Lower glume - Beak tip:LongLower glume - Shoulder width:MediumLower glume - Shoulder type:Elevated

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Strong

Lower glume - Internal imprint:

Lower lemma - Beak shape:Moderately curved to strongly curved

Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:*Grain - colour:

Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: Light to medium

*Seasonal type: Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 28 March 2017



MARSTON

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature strongly curved to geniculate

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol light to medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence very late



Variety Name: Maurizio crop: Winter Wheat

Breeder's Reference: SZD 6374 AFP No:

Breeder: Saatzucht Donau GmbH Translated from: Hungarian

Origin: Date: 19 November 2018

Code: IZIO

Coleoptile-anthocyanin colouration:Abesnt or very weak*Plant - growth habit:Intermediate*Time ear emergence:Early to medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Very low to low.

*Flag leaf - Glaucosity of sheath: Strong

Flag leaf - Glaucosity of leaf blade: Medium to strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Very strong to strong

Hairiness of uppermost node:

*Plant - height:

*Straw - wall thickness @ 20cms below ear: Thin
*Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White *Ear - Shape (side profile): Parallel

Ear - Length:Short to medium*Ear - Density:Lax to medium*Awns or scurs - presence:Awns present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Long

*Awns at tip of ear - length:

Lower glume - Beak length: Short

Lower glume - Beak shape (curvature): Moderately curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Medium

Lower glume - Shoulder type: Sloping to sightly sloping

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: Medium to dark

*Seasonal type: Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 19 March 2019



MAXWELL

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presenceawns presentAwns or scurs - distributionwhole lengthAwns or scurs - lengthvery long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length very long

Beak curvature moderate to strongly curved

Shoulder width narrow

Shoulder shape strongly elevated with 2nd pt. present

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium
Wall thickness @20cms below ear very thick or filled

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) weak to medium

Culm - neck medium

Ear very weak to weak

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence very early



MAYFLOWER

CHARACTERS IN DETAIL

Ear

Colour white

short to medium Length

Density dense Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface

short to medium Beak length

Beak curvature moderate to strongly curved

Shoulder width narrow to medium Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) weak to medium

Culm - neck strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong to very strong Plant - growth habit intermediate Time of ear emergence medium to late Flag leaf - anthocyanin of auricles absent or very weak



MEMPHIS

CHARACTERS IN DETAIL

Grain

Colour reddish
Colouration with phenol medium

Plant

Anthocyanin colouration of coleoptile very weak to weak

Growth habit intermed. to semi-prostrate

Frequency of plants with recurved flag leaves lo

Anthocyanin colouration of flag leaf auricles absent or very weak Time of ear emergence medium to late Length (stem, ears, awns or scurs) very short to short

Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong

Flag leaf blade (lower side) strong
Ear strong
Culm - neck strong

Lower glume

Hairiness of external surface absent
Shoulder width narrow

Shoulder shape slightly elevated Length of beak short to medium

Shape of beak slight to moderately curved Extent of internal hairs very large (Group 3)

External surface roughness rough

Ear

Density medium to dense
Length (excluding awns or scurs) short to medium
Awns or scurs - presence scurs present
Awns or scurs - length medium to long

Colour white

Shape in profile parallel sided
Apical rachis hairiness on convex surface absent or very small

Awns or scurs - distribution upper 1/2



MERIT

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium (group 1+)

Beak length lor

Beak curvature moderate to strongly curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



MINDFUL

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium
Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium (group 1+)

Beak length short

Beak curvature strongly curved Shoulder width medium

Shoulder shape horizontal to slightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



MONROE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlongApical rachis segment - convex face hairinessmedium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence late



MONTEREY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ¾

Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



Variety Name: **MORTIMER** crop: Winter Wheat

Breeder's Reference: SC 2405 AFP No:

Breeder: Secobra Translated from:

Origin: 15 May 2018

Code: MORT

Coleoptile-anthocyanin colouration:Absent or very weak*Plant - growth habit:Semi prostrate*Time ear emergence:Medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Very low ot low

*Flag leaf - Glaucosity of sheath: Strong

Flag leaf - Glaucosity of leaf blade: Strong to very strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong

Hairiness of uppermost node:

*Plant - height: Short

*Straw - wall thickness @ 20cms below ear: Absent or very thin *Ear - Glaucosity: Medium to strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White
*Ear - Shape (side profile): Parallel
Ear - Length: Long

*Ear - Density: Lax to medium
*Awns or scurs - presence: Present

Awns or scurs - distibution: *Scurs at tip of ear - length:

*Awns at tip of ear - length: Short to medium

Lower glume - Beak length: Short

Lower glume - Beak shape (curvature):Straight to slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow to medium

Lower glume - Shoulder type:

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Very weak to weak

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Weak to medium

*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:*Seasonal type:

Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 16 May 2018



MOSAIC

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 3/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (3)
Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect to intermediate
Time of ear emergence late to very late



Variety Name: Moschus crop: Winter wheat

Breeder's Reference: STRU 114017 AFP No:

Breeder: Dr HermannStrube **Translated from:** German

Origin: Date: 19 November 2018

Code: MOSC

Coleoptile-anthocyanin colouration: Strong to very strong

*Plant - growth habit: Intermediate
*Time ear emergence: Medium to late

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves:Low to medium*Flag leaf - Glaucosity of sheath:Strong to very strongFlag leaf - Glaucosity of leaf blade:Strong to very strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong to very strong

Hairiness of uppermost node:

*Plant - height: Medium

*Straw - wall thickness @ 20cms below ear: Absent or very thin

*Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour:

*Ear - Shape (side profile): Tapering
Ear - Length: Medium to long
*Ear - Density: Lax to medium
*Awns or scurs - presence: Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short to short

*Awns at tip of ear - length:

Lower glume - Beak length: Short

Lower glume - Beak shape (curvature): Slightly to moderately curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Narrow to medium

Lower glume - Shoulder type: Straight

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Medium to strong

*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:*Seasonal type:

Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 19 March 2019



MOULTON

CHARACTERS IN DETAIL

Ear

white Colour Length long Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)
Beak length short to medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence late to very late



MV FREDERICIA

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profiletaperingAwns or scurs - presenceawns presentAwns or scurs - distributionwhole lengthAwns or scurs - lengthvery long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length long to very long

Beak curvature moderate to strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



MYRIAD

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ¾
Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



MYSTIC CS

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long

Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slight to moderately curved Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence medium to late



NELSON

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper ½ Awns or scurs - length: medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape straight

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



OAKLEY

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ¾
Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) medium
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium



OLIVIN

CHARACTERS IN DETAIL

Ear

Colour: white

Length: medium

Density: medium

Shape (side profile) parallel sided

Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short

Apical rachis segment - convex face hairiness: medium to strong

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: medium - group 2

Beak length: very short to short

Beak shape (curvature): strongly curved

Shoulder narrow to medium

Shoulder slightly sloping to straight

Lower Lemma

Beak shape: strongly curved to geniculate

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: medium to long

Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong
Flag leaf - blade (lower side): medium

Culm neck: medium

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin strong to very strong

Plant - growth habit: intermediate to semi-prostrate

Time ear emergence: late

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



OLYMPUS

long to very long

CHARACTERS IN DETAIL

Awns or scurs - length

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 3/4

Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



Variety Name: Opus crop: Winter Wheat

Breeder's Reference: PBIS 99/70 **AFP No:**

Breeder: Erhardt Eger Translated from: German

Origin: Date: 23 November 2001

Code: OPUS

Coleoptile-anthocyanin colouration: absent or very weak

*Plant - growth habit: intermediate to semi-prostrate

medium

*Time ear emergence: medium to late
Flag leaf - Anthocyanin colouration of auricles: absent or very weak
Frequency of plants with recurved flag leaves: very low to low

*Flag leaf - Glaucosity of sheath: strong

Flag leaf - Glaucosity of leaf blade: medium to strong

Anthers - anthocyanin colouration:

Glaucosity of neck: strong

Hairiness of uppermost node:
*Plant - height:

*Straw - wall thickness @ 20cms below ear: very thin *Ear - Glaucosity: medium

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: white

*Ear - Shape (side profile): tapering

Ear - Length: medium

*Ear - Density: medium

*Ear - Density: medium

*Awns or scurs - presence: scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length:
short

*Awns at tip of ear - length: Lower glume - Beak length:

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width: Lower glume - Shoulder type:

Lower glume - External surface roughness: Lower glume - Extent of internal hairs:

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour: Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type: winter

Extra characters:

* = Characteristics which should always be included in the description of the variety.



06 May 2020

ORATOR

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs present

Awns or scurs - distribution upper 1/4
Awns or scurs - length medium
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)
Beak length medium to long

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) weak to medium
Culm - neck medium to strong
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit semi-prostrate
Time of ear emergence medium to late



ORBIT

CHARACTERS IN DETAIL

Ear

white Colour Length long Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length short to medium Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium

Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



OSMIUM

CHARACTERS IN DETAIL

Ear

Colour white
Length short
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface rough
Extent of internal hairs weak (group 1)

Beak length short

Beak curvature v.slightly curved Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



OXFORD

CHARACTERS IN DETAIL

Ear

Colour white
Length medium to long
Density medium to dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long to very long

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium

Beak curvaturemoderately curvedShoulder widthnarrow to mediumShoulder shapeslightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish
Colouration with phenol dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-prostrate

Time of ear emergence late to very late

Flag leaf - anthocyanin of auricles absent or very weak



PANACEA

absent

CHARACTERS IN DETAIL

Anthers - anthocyanin colouration

Ear

Colour white

Length medium to long
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ¾
Awns or scurs - length: long
Apical rachis segment - convex face hairiness medium

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late



PANORAMA

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper ¼

Awns or scurs - length: short to medium
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) weak to medium

Culm - neck medium

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



PHILOSOPHER

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ¾
Awns or scurs - length: long
Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



PIRENEO

CHARACTERS IN DETAIL

Ear

Colour: white Length: medium

Density: medium to dense

Shape (side profile) tapering
Awns or scurs - presence: awns present

Awns or scurs - distribution:

Awns or scurs - length: long

Apical rachis segment - convex face hairiness: medium

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: medium - group 2

Beak length: long to very long

Beak shape (curvature): slightly curved

Shoulder narrow Shoulder straight

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: medium

Straw

Length: medium Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: weak

Flag leaf - blade (lower side): very weak to weak

Culm neck: weak to medium

Ear - Glaucosity: weak

Plant growth characters

Coleoptile-anthocyanin

Plant - growth habit: intermediate
Time ear emergence: early to late

Flag leaf - Anthocyanin colouration of auricles:



Variety Name: **PONTICUS** crop: Winter Wheat

Breeder's Reference: STRU 114010 AFP No:

Breeder: Dr Hermann Strube **Translated from:** German

Origin: Date: 14 November 2017

Code: PONT

Coleoptile-anthocyanin colouration: Strong

*Plant - growth habit: Intermediate to semi-prostrate

*Time ear emergence: Medium to late

Flag leaf - Anthocyanin colouration of auricles:

*Flag leaf - Glaucosity of leaf blade:

Very low to low

Very strong

Very strong

Very strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Very strong

Hairiness of uppermost node:

*Plant - height: Strong to medium
*Straw - wall thickness @ 20cms below ear: Absent or very thin

*Ear - Glaucosity: Strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Tapering

Ear - Length: Medium to long

*Ear - Density: Lax

*Awns or scurs - presence: Scurs present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Very short to short

*Awns at tip of ear - length:

Lower glume - Beak length: Very short to short

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width:Narrow to mediumLower glume - Shoulder type:Slightly sloping

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Weak

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour:

Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:*Seasonal type:

Winter

Extra characters:

^{* =} Characteristics which should always be included in the description of the variety. 28 March 2018



Variety Name: PRAKTIK Crop:

Breeder's Reference: AFP No: 1/

Breeder: RAGT Translated from:

Origin: GERMANY Date: 22 October 2012

Code: PRAK

Coleoptile-anthocyanin colouration: ABSENT OR VERY WEAK

*Plant - growth habit: INTERMEDIATE TO SEMI-PROSTRATE

*Time ear emergence: MEDIUM TO LATE

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: ABSENT OR VERY LOW

*Flag leaf - Glaucosity of sheath: MEDIUM
Flag leaf - Glaucosity of leaf blade: MEDIUM

Anthers - anthocyanin colouration:

Glaucosity of neck: MEDIUM TO STRONG

Hairiness of uppermost node:

*Plant - height: SHORT

*Straw - wall thickness @ 20cms below ear: very thin

*Ear - Glaucosity: MEDIUM

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: WHITE
*Ear - Shape (side profile): TAPERING

Ear - Length: SHORT TO MEDIUM

*Ear - Density: MEDIUM

*Awns or scurs - presence: SCURS PRESENT

Awns or scurs - distibution:

*Scurs at tip of ear - length: VERY SHORT TO SHORT

*Awns at tip of ear - length:

Lower glume - Beak length: VERY SHORT TO SHORT

Lower glume - Beak shape (curvature): STRAIGHT

Lower glume - Beak tip:SLIGHTLY CURVEDLower glume - Shoulder width:NARROW TO MEDIUM

Lower glume - Shoulder type:

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: MEDIUM

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: MEDIUM *Grain - colour: RED

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: DARK

*Seasonal type: WINTER TYPE

Extra characters: STRAW:PILTH IN CROSS SECTION -

VERY THIN

^{* =} Characteristics which should always be included in the description of the variety. 30 March 2015



QI

medium

CHARACTERS IN DETAIL

Awns or scurs - length:

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ¼

Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong - Group 2+

Beak lengthmediumBeak curvatureslightly curvedShoulder widthnarrowShoulder shapeelevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit semi-erect
Time of ear emergence medium



RA3W 08

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium
Density: medium to dense

Shape (side profile) tapering
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness: weak to medium

Anthers - anthocyanin

Lower Glume

External surface roughness: medium

Extent of internal hairs: medium - group 2

Beak length: medium

Beak shape (curvature): moderately curved

Shoulder broad
Shoulder elevated

Lower Lemma

Beak shape: moderately curved to strongly curved

Grain

Colour red

Colouration with phenol: medium to dark

Straw

Length: short to medium

Wall thickness @ 20cms below ear: thin

Glaucosity

Flag leaf - sheath: strong to very strong Flag leaf - blade (lower side): medium to strong

Culm neck: weak
Ear - Glaucosity: medium

Plant growth characters

Coleoptile-anthocyanin medium
Plant - growth habit: intermediate
Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles:



RADAR

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol medium to dark

Plant

Anthocyanin colouration of coleoptile weak

Growth habit intermediate
Frequency of plants with recurved flag leaves very low to low
Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late
Length (stem, ears, awns or scurs) short
Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong

Flag leaf blade (lower side) medium to strong
Ear weak to medium
Culm - neck medium to strong

Lower glume

Hairiness of external surface absent Shoulder width narrow

Shoulder shape slightly elevated
Length of beak medium to long
Shape of beak strongly curved
Extent of internal hairs very large (Group 3)
External surface roughness slightly rough

Ear

Density dense

Length (excluding awns or scurs)short to mediumAwns or scurs - presencescurs presentAwns or scurs - lengthlong to very longColourwhite

Shape in profile parallel sided
Apical rachis hairiness on convex surface small to medium
Awns or scurs - distribution upper 1/2



RAMSAY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)

Beak length short

Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



RAPSODIA

CHARACTERS IN DETAIL

Ear

Colour: white

Length: short to medium

Density: medium

Shape (side profile) tapering

Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: short

Apical rachis segment - convex face hairiness: weak to medium

Anthers - anthocyanin

Lower Glume

External surface roughness: smooth

Extent of internal hairs: medium to strong - group 2 to 3

Beak length: short

Beak shape (curvature): moderately curved

Shoulder medium Shoulder straight

Lower Lemma

Beak shape: slightly curved

Grain

Colour red
Colouration with phenol: medium

Straw

Length: short

Wall thickness @ 20cms below ear: very thin to thin

Glaucosity

Flag leaf - sheath: strong to very strong medium to strong

Culm neck: medium to strong

Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin strong

Plant - growth habit: semi-prostrate to prostrate

Time ear emergence: medium

Flag leaf - Anthocyanin colouration of auricles: absent or very weak



RAZZANO

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs – distribution

Awns or scurs - length: very short

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface

Extent of internal hairs medium to strong - Group 2+

Beak length short

Beak curvature v.slightly curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



REFLECTION

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length medium
Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs strong
Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence medium to late



RELAY

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium to dense

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns or scurs - distribution upper ½

Awns or scurs - length: long

Awns or scurs - length: long
Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



REVELATION

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ¼
Awns or scurs - length: long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



RGT ADVENTURE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length short to medium
Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles

(NIAB

RGT ANELLE

CHARACTERS IN DETAIL

Ear

Colour coloured
Length short to medium

Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length very short to sl

Awns or scurs - length very short to short
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) medium to strong
Culm - neck strong to very strong
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



RGT BAIRSTOW

absent

CHARACTERS IN DETAIL

Anthers - anthocyanin colouration

Ear

Colour white

Length short to medium
Density medium

Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4

Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness very strong

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



RGT BAKEWELL

medium to dense

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol medium to dark

Plant

Anthocyanin colouration of coleoptile strong

Growth habit intermed. to semi-prostrate

Frequency of plants with recurved flag leaves absent or very low Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late Length (stem, ears, awns or scurs) short Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong Flag leaf blade (lower side) strong to very strong

strong to very strong Culm - neck strong to very strong

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium Shoulder shape slightly elevated Length of beak short to medium Shape of beak moderately curved Extent of internal hairs very large (Group 3)

External surface roughness smooth

Ear

Density Length (excluding awns or scurs) short to medium Awns or scurs - presence scurs present

Awns or scurs - length very short to short

Colour white

Shape in profile parallel sided Apical rachis hairiness on convex surface medium to large

Awns or scurs - distribution upper 1/4



RGT BLOSSOM

weak to medium

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak lengthshort to mediumBeak curvaturemoderately curvedShoulder widthnarrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence late



RGT BOWEN

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length long
Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence late



RGT COOK

very long

CHARACTERS IN DETAIL

Awns or scurs - length

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presenceawns presentAwns or scurs - distributionwhole length

Apical rachis segment - convex face hairiness strong to very strong

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough Extent of internal hairs strong to whole surface

Beak length long to very long
Beak curvature strongly curved

Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Straw

Colour red
Colouration with phenol dark

Length very short to short Wall thickness @20cms below ear absent or very thin

•

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



RGT CONVERSION

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long to very long

Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth Extent of internal hairs medium Beak length long

Beak curvature strongly curved Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red Colouration with phenol dark

Straw

short Length

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium Culm - neck medium

weak to medium Ear

Plant growth characters

absent to weak Coleoptile - anthocyanin colouration Plant - growth habit intermediate late

Time of ear emergence



RGT DALLAGLIO

absent

CHARACTERS IN DETAIL

Anthers - anthocyanin colouration

Ear

white Colour Length medium to long Density medium to dense Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness large

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak lengthmediumBeak curvaturestrongly curvedShoulder widthnarrow to mediumShoulder shapeslightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish
Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong

Ear Strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak

Plant - growth habit semi-prostrate

Time of ear emergence late to very late

Flag leaf - anthocyanin of auricles absent or very weak



RGT DOMINICI

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium
Apical rachis segment - convex face hairiness large
Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs medium (group 2)

Beak lengthmediumBeak curvaturestrongly curvedShoulder widthmedium to broadShoulder shapeslightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



RGT DRAX

CHARACTERS IN DETAIL

Ear

white Colour medium Length Density dense Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak lengthmediumBeak curvatureslightly curvedShoulder widthnarrow to mediumShoulder shapestraight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence

Flag leaf - anthocyanin of auricles absent or very weak



RGT EVESHAM

CHARACTERS IN DETAIL

Grain

Colour reddish Colouration with phenol dark

Plant

Anthocyanin colouration of coleoptile medium

Growth habit intermed. to semi-prostrate

Frequency of plants with recurved flag leaves medium

Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late
Length (stem, ears, awns or scurs) short
Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong
Flag leaf blade (lower side) strong

Ear medium to strong

Culm - neck strong

Lower glume

Hairiness of external surface absent
Shoulder width narrow

Shoulder shape slightly elevated Length of beak medium

Shape of beak moderately curved Extent of internal hairs medium (Group 2)

External surface roughness entire external surface rough

Ear

Density dense

Length (excluding awns or scurs)short to mediumAwns or scurs - presencescurs presentAwns or scurs - lengthmediumColourwhiteShape in profileparallel sided

Apical rachis hairiness on convex surface very small to small

Awns or scurs - distribution upper 1/2



RGT FLINTOFF

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence awns present
Awns or scurs - distribution whole length
Awns or scurs - length very long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough
Extent of internal hairs whole internal surface covered

Beak length very long

Beak curvature moderate to strongly curved Shoulder width very narrow to narrow

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence medium to late

Flag leaf - anthocyanin of auricles absent or very weak



RGT GALACTUS

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate



Variety Name: RGT GRADIENT Crop: WW

Breeder's Reference: RW41394 AFP No:

Breeder: R2n sas **Translated from:** Danish

Origin: Trident + Gravitas Date: 22 March 2016

Code: RGTG

Coleoptile-anthocyanin colouration: Very weak to weak

*Plant - growth habit: Prostrate

*Time ear emergence: Medium

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Low to medium

*Flag leaf - Glaucosity of sheath: Strong

Flag leaf - Glaucosity of leaf blade: Medium to strong

Anthers - anthocyanin colouration:

Glaucosity of neck: Strong

Hairiness of uppermost node:

*Plant - height: Short to medium

*Straw - wall thickness @ 20cms below ear: Absent or Very thin

*Ear - Glaucosity: Medium to strong

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White *Ear - Shape (side profile): Tapering

Ear - Length: Short to medium

*Ear - Density: Medium
*Awns or scurs - presence: Present

Awns or scurs - distibution:

*Scurs at tip of ear - length: Long

*Awns at tip of ear - length:

Lower glume - Beak length:Medium to longLower glume - Beak shape (curvature):Slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Medium **Lower glume - Shoulder type:** Elevated

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Medium

Lower glume - Internal imprint:

Lower lemma - Beak shape: Moderately curved

Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: Weak to medium

*Grain - colour: Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol: *Seasonal type:Dark
Winter

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 28 March 2018

RGT GRAVITY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profiletaperingAwns or scurs - presencescurs present

Awns or scurs - distribution upper 1/2
Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Glaucosity

Length short to medium Wall thickness @20cms below ear absent or very thin

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



RGT GROUSE

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long
Apical rachis segment - convex face hairiness large
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium
Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong

Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



RGT HEXTON

CHARACTERS IN DETAIL

Grain

Colour reddish
Colouration with phenol medium

Plant

Anthocyanin colouration of coleoptile very weak to weak

Growth habit intermed. to semi-prostrate

Frequency of plants with recurved flag leaves very low to low

Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late
Length (stem, ears, awns or scurs) short
Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong

Flag leaf blade (lower side) medium to strong
Ear medium to strong

Culm - neck strong

Lower glume

Hairiness of external surface absent
Shoulder width medium

Shoulder shape slightly elevated Length of beak medium

Shape of beak moderate to strongly curved Extent of internal hairs very small to medium (Group 1+)

External surface roughness smooth

Ear

Density medium to dense
Length (excluding awns or scurs) medium to long
Awns or scurs - presence scurs present

Awns or scurs - length medium Colour white

Shape in profile parallel sided
Apical rachis hairiness on convex surface medium to large

Awns or scurs - distribution upper 1/2



RGT HIGHGROVE

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profiletaperingAwns or scurs - presenceawns presentAwns or scurs - distributionwhole lengthAwns or scurs - lengthvery long

Apical rachis segment - convex face hairiness very small to small

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length very long

Beak curvature moderate to strongly curved Shoulder width very narrow to narrow

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour reddish Colouration with phenol medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



RGT HOWLEY

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness large to very large

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium to long
Beak curvature moderately curved
Shoulder width narrow to medium

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-prostrate

Time of ear emergence late to very late

Flag leaf - anthocyanin of auricles absent or very weak



RGT ILLUSTRIOUS

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length medium Apical rachis segment - convex face hairiness strong Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs medium to strong (2+)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late



RGT KNIGHTSBRIDGE

CHARACTERS IN DETAIL

Ear

Colour white medium Length medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) weak
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence very late



RGT KOI

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness strong
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length medium
Beak curvature strongly curved
Shoulder width narrow

Shoulder shape slightly sloping

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong

Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate



RGT LANTERN

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense

Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length long to very long

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late



RGT LUDLOW

CHARACTERS IN DETAIL

Grain

Colour reddish
Colouration with phenol dark

Plant

Anthocyanin colouration of coleoptile weak

Growth habit intermed. to semi-prostrate

Frequency of plants with recurved flag leaves very low to low
Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late
Length (stem, ears, awns or scurs) short
Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong

Flag leaf blade (lower side) medium to strong

Ear strong
Culm - neck strong

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium

Shoulder shape horizontal to slightly elevated

Length of beak medium

Shape of beak slight to moderately curved Extent of internal hairs very small to medium (Group 1+)

External surface roughness smooth

Ear

Density dense
Length (excluding awns or scurs) medium
Awns or scurs - presence scurs present
Awns or scurs - length short to medium

Colour white

Shape in profile parallel sided
Apical rachis hairiness on convex surface large to very large

Awns or scurs - distribution upper 1/4



RGT MARLBOROUGH

medium

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlong

Anthers - anthocyanin colouration

Apical rachis segment - convex face hairiness

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)
Beak length medium to long
Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence late to very late



RGT MOGUL

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long

Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved
Shoulder width very narrow to narrow
Shoulder shape slightly elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence

Flag leaf - anthocyanin of auricles absent or very weak



RGT NONU

medium to large

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Anthers - anthocyanin colouration absent

Apical rachis segment - convex face hairiness

Lower glume

External surface rough

Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature strongly curved

Shoulder width narrow to medium

Shoulder shape horizontal to slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol dark to very dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak

Plant - growth habit semi-prostrate

Time of ear emergence late to very late

Flag leaf - anthocyanin of auricles absent or very weak



RGT PADDINGTON

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness medium

A ...

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) weak to medium

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence very late



RGT PARISSE

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length very long Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium Beak curvature strongly curved Shoulder width narrow to medium Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong Flag leaf- blade (lower side) strong Culm - neck strong Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-prostrate Time of ear emergence late to very late Flag leaf - anthocyanin of auricles absent or very weak



RGT PEMBROKE

long

CHARACTERS IN DETAIL

Awns or scurs - length

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence late to very late



RGT QUICKSILVER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate



RGT RASHID

CHARACTERS IN DETAIL

Ear

white Colour medium Length Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence very late

Flag leaf - anthocyanin of auricles absent or very weak



RGT SACRAMENTO

CHARACTERS IN DETAIL

Ear

white Colour Length short Density medium Shape-side profile tapering Awns or scurs - presence awns present Awns or scurs - distribution whole length Awns or scurs - length very long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length long to very long
Beak curvature slightly curved
Shoulder width very narrow to narrow
Shoulder shape slightly sloping

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol light

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong Ear strong

Plant growth characters

Coleoptile - anthocyanin colourationstrongPlant - growth habitsemi-erectTime of ear emergencemedium to late



RGT SAKI

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length long

Beak curvature strongly curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate



RGT SILVERSURFER

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length long

Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate



RGT SCRUMMAGE

CHARACTERS IN DETAIL

Ear

Colour white
Length medium to long
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/2
Awns or scurs - length medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence late



RGT SORCERER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymediumShape-side profiletaperingAwns or scurs - presenceawns present

Awns or scurs - distribution

Awns or scurs - length n/a

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length long

Beak curvature moderate to strongly curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence very late



RGT STOKES

medium to long

CHARACTERS IN DETAIL

Awns or scurs - length

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermediate

Time of ear emergence late to very late

Flag leaf - anthocyanin of auricles absent or very weak



RGT TINDALL

CHARACTERS IN DETAIL

Ear

Colour white

Length long to very long Density lax to medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length long

Beak curvature strongly curved

Shoulder width narrow

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-prostrate Time of ear emergence late to very late Flag leaf - anthocyanin of auricles absent or very weak



RGT UNIVERSE

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



RGT VENOM

CHARACTERS IN DETAIL

Ear

Colour white Length long

Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate



RGT WASABI

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium
Density dense
Shape-side profile parallel sided

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate



RGT WESTMINSTER

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present

Awns or scurs - distribution

Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) weak to medium

Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium Plant - growth habit intermediate Time of ear emergence very late



RGT WILKINSON

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs medium (group 2)
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-prostrate
Time of ear emergence late to very late
Flag leaf - anthocyanin of auricles absent or very weak



RGT WOLVERINE

CHARACTERS IN DETAIL

Ear

white Colour medium Length Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length short Apical rachis segment - convex face hairiness strong Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)

Beak length short

Beak curvature slightly curved Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong

Plant - growth habit intermed. to semi-prostrate



RGT ZINZAN

CHARACTERS IN DETAIL

Ear

Colour white
Length medium to long
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness small to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough

Extent of internal hairs strong (group 3)
Beak length short to medium

Beak curvature moderate to strongly curved

Shoulder width narrow

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



RIBAND

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs weak (1)
Beak length medium

Beak curvature slight to moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) weak
Culm - neck medium
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak

Plant - growth habit prostrate
Time of ear emergence medium

Flag leaf - anthocyanin of auricles absent or very weak



RILEY

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol medium to dark

Plant

Anthocyanin colouration of coleoptile very weak to weak

Growth habit intermed. to semi-prostrate

Frequency of plants with recurved flag leaves

Anthocyanin colouration of flag leaf auricles

Time of ear emergence

Length (stem, ears, awns or scurs)

very low to low
absent or very weak
medium to late
very short

Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong

Flag leaf blade (lower side) strong
Ear strong
Culm - neck strong

Lower glume

Hairiness of external surface absent
Shoulder width narrow

Shoulder shape horizontal to slightly elevated

Length of beak medium

Shape of beak slight to moderately curved Extent of internal hairs very large (Group 3)

External surface roughness smooth

Ear

Density dense

Length (excluding awns or scurs)short to mediumAwns or scurs - presencescurs presentAwns or scurs - lengthlong

Colour white

Shape in profile parallel sided
Awns or scurs - distribution upper 1/2



ROADSTER

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence awns present

Awns or scurs - distribution

Awns or scurs - length n/a
Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length long to very long
Beak curvature moderately curved
Shoulder width very narrow to narrow

Shoulder shape straight

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol light

Straw

Length very short to short
Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



ROBIGUS

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper ¼

Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2
Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate
Time of ear emergence medium
Flag leaf - anthocyanin of auricles weak



ROMA

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol medium to dark

Plant

Anthocyanin colouration of coleoptile weak to medium

Growth habit intermed. to semi-prostrate

Frequency of plants with recurved flag leaves absent or very low Anthocyanin colouration of flag leaf auricles absent or very weak Time of ear emergence medium to late

Length (stem, ears, awns or scurs) very short to short

Straw wall thickness (20cm below ear) th

Glaucosity

Flag leaf sheath strong to very strong Flag leaf blade (lower side) medium to strong

Ear strong
Culm - neck strong

Lower glume

Hairiness of external surface absent Shoulder width narrow

Shoulder shape Slightly elevated Length of beak short to medium

Shape of beak moderate to strongly curved

Extent of internal hairs very large (Group 3)

External surface roughness entire external surface rough

Ear

Density medium
Length (excluding awns or scurs) medium

Awns or scurs - presence scurs present

Awns or scurs - length long
Colour white

Shape in profile parallel sided
Apical rachis hairiness on convex surface small to medium

Awns or scurs - distribution upper 3/4



ROWLING

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length short to medium
Beak curvature slightly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



ROYSTON

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)
Beak length short to medium
Beak curvature strongly curved
Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) weak to medium
Culm - neck weak to medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate
Time of ear emergence late
Flag leaf - anthocyanin of auricles



RUFUS

CHARACTERS IN DETAIL

Grain

Colour reddish Colouration with phenol dark

Plant

Anthocyanin colouration of coleoptile weak
Growth habit intermediate

Frequency of plants with recurved flag leaves low

Anthocyanin colouration of flag leaf auricles absent or very weak Time of ear emergence medium to late

Length (stem, ears, awns or scurs) short Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong
Flag leaf blade (lower side) strong

Ear very weak to weak Culm - neck weak to medium

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium

Shoulder shape horizontal to slightly elevated

Length of beak medium
Shape of beak strongly curved
Extent of internal hairs medium (Group 2)
External surface roughness slightly rough

Ear

Density dense

Length (excluding awns or scurs)medium to longAwns or scurs - presencescurs presentAwns or scurs - lengthvery longColourwhite

Shape in profile parallel sided
Awns or scurs - distribution upper 3/4



RUSKIN

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense Shape-side profile tapering

Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium
Beak length short to medium
Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



SAHARA

CHARACTERS IN DETAIL

Ear

Colour white

Length medium

Density lax to medium

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns or scurs - distribution upper ½

Awns or scurs - length: short to medium
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length medium to long

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium
Flag leaf - blade (lower side) weak
Culm - neck weak

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late to very late



SARTORIAL

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthlongApical rachis segment - convex face hairinessstrong

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs weak (group 1)
Beak length medium to long
Beak curvature moderately curved

Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermediate
Time of ear emergence late



SAVELLO

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong (group 3)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence very late



SCOUT

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length: medium Apical rachis segment - convex face hairiness weak Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature straight to slightly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



SCRIBE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ¼
Awns or scurs - length: medium
Apical rachis segment - convex face hairiness strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature slightly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate
Time of ear emergence medium



SHABRAS

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthvery long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong (group 3)
Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape strongly elevated with 2nd pt. present

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



SHEPHERD

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium (2)

Beak length medium
Beak curvature moderat

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong
Plant - growth habit intermediate
Time of ear emergence medium to late



SHERLOCK

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium
Shape-side profile tapering
Awns or scurs - presence awns present
Awns or scurs - distribution whole length
Awns or scurs - length very long

Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth Extent of internal hairs medium (2)

Beak length long to very long

Beak curvature moderate to strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



SHIRE

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length long
Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (1+)

Beak length medium

Beak curvature slightly curved

Shoulder width narrow

Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



SKYFALL

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile parallel sided Awns or scurs - presence awns present Awns or scurs - distribution whole length Awns or scurs - length: very long Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3 Beak length long to very long Beak curvature v.slightly curved Shoulder width very narrow to narrow Shoulder shape elev. to elev with 2nd point

Lower lemma

Beak curvature straight

Grain

Colour red

Colouration with phenol dark to very dark

Straw

short to medium Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong Ear

strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium



SMUGGLER

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile tapering Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium Apical rachis segment - convex face hairiness medium Anthers - anthocyanin colouration absent

Lower glume

External surface entire external surface rough Extent of internal hairs weak to medium (group 1+)

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration very strong
Plant - growth habit intermediate

Time of ear emergence early

Flag leaf - anthocyanin of auricles absent or very weak



SOISSONS

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profiletaperingAwns or scurs - presenceawns presentAwns or scurs - distributionwhole length

Awns or scurs - length: long

Apical rachis segment - convex face hairiness absent to weak

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs weak - Group 1

Beak length Beak curvature

Shoulder width narrow

Shoulder shape slightly sloping

Lower lemma

Beak curvature straight

Grain

Colour red

Colouration with phenol

Straw

Length

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath absent or very weak

Flag leaf- blade (lower side) weak

Culm - neck absent or very weak Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate

Time of ear emergence very early to early Flag leaf - anthocyanin of auricles absent or very weak



SOLACE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ½
Awns or scurs - length: long

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width medium Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



SOLSTICE

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density lax to medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ½
Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape straight

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side)

Culm - neck strong to very strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate

Time of ear emergence late
Flag leaf - anthocyanin of auricles very weak



SOLTEK

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density dense
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length short to medium

Aprical rachis segment - convex face hairiness strong to very strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)
Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



Variety Name: **SORRIAL** Crop: WINTER WHEAT

Breeder's Reference: AFP No:

Breeder: CAUSSADE SEMENCES SA Translated from:

Origin: FRANCE Date: 03 June 2013

Code: SORRI

Coleoptile-anthocyanin colouration: MEDIUM

*Plant - growth habit: INTERMEDIATE TO SEMI-PROSTRATE

*Time ear emergence: EARLY TO MEDIUM

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: *Flag leaf - Glaucosity of sheath:VERY LOW TO LOW
VERY STRONG

Flag leaf - Glaucosity of leaf blade: STRONG TO VERY STRONG

Anthers - anthocyanin colouration:

Glaucosity of neck: STRONG TO VERY STRONG

Hairiness of uppermost node:

*Plant - height: SHORT

*Straw - wall thickness @ 20cms below ear: VERY THIN

*Ear - Glaucosity: STRONG TO VERY STRONG

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: WHITE

*Ear - Shape (side profile): PARALLEL SIDED
Ear - Length: MEDIUM TO LONG

*Ear - Density: DENSE

*Awns or scurs - presence: SCURS PRESENT

Awns or scurs - distibution:

*Scurs at tip of ear - length: VERY SHORT TO SHORT

*Awns at tip of ear - length: Lower glume - Beak length:

Lower glume - Beak shape (curvature):

Lower glume - Beak tip:

Lower glume - Shoulder width: MEDIUM TO BROAD

Lower glume - Shoulder type: STRAIGHT

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: WEAK

Lower glume - Internal imprint:

Lower lemma - Beak shape: SLIGHTLY CURVED

Lower lemma - Beak length-: SHORT

Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness: WEAK TO MEDIUM

*Grain - colour: RED

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type:

WINTER

Extra characters:



^{* =} Characteristics which should always be included in the description of the variety. 30 March 2015

SPEARHEAD

medium to strong

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium to long

Anthers - anthocyanin colouration absent

Apical rachis segment - convex face hairiness

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface
Beak length short to medium
Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong
Flag leaf- blade (lower side) very strong
Culm - neck very strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



SPOCK

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present upper 1/4 Awns or scurs - distribution Awns or scurs - length medium to long Apical rachis segment - convex face hairiness strong to very strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence late to very late



SPYDER

medium to long

CHARACTERS IN DETAIL

Awns or scurs - length

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Apical rachis segment - convex face hairiness weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth Extent of internal hairs medium (2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



STIGG

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density very dense

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns or scurs - distribution upper ¼

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved
Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) strong to very strong

Culm - neck strong

Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



STRATOSPHERE

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense
Shape-side profile parallel sided

Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length very short to short Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium

Flag leaf- blade (lower side) weak

Culm - neck medium

Culm - neck medium
Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late to very late



SWALLOW

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long to very long Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs medium to strong (group 2+)

Beak length long

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol medium

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong

Culm - neck strong to very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



SY CHEER

CHARACTERS IN DETAIL

Grain

Colour reddish

Colouration with phenol medium to dark

Plant

Anthocyanin colouration of coleoptile weak

Growth habit intermed. to semi-prostrate

Frequency of plants with recurved flag leaves very low to low Anthocyanin colouration of flag leaf auricles absent or very weak Time of ear emergence medium to late

short Length (stem, ears, awns or scurs)

Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong

Flag leaf blade (lower side) strong to very strong

Ear strong

Culm - neck strong to very strong

Lower glume

Hairiness of external surface absent

Shoulder width narrow to medium Shoulder shape slightly elevated medium

Length of beak

Shape of beak moderately curved Extent of internal hairs medium (Group 2)

External surface roughness smooth

Ear

Density medium to dense

Length (excluding awns or scurs) medium Awns or scurs - presence scurs present Awns or scurs - length medium to long

Colour white

Shape in profile parallel sided

Apical rachis hairiness on convex surface small

Awns or scurs - distribution upper 1/4



SY COACH

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4

Awns or scurs - length very short to short
Apical rachis segment - convex face hairiness large to very large

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium
Beak curvature strongly curved
Shoulder width medium

Shoulder shape slightly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late
Flag leaf - anthocyanin of auricles absent or very weak



SY EPSON

CHARACTERS IN DETAIL

Ear

Colour: white Length: long

Density: lax to medium
Shape (side profile) parallel sided
Awns or scurs - presence: scurs present

Awns or scurs - distribution:

Awns or scurs - length: long

Apical rachis segment - convex face hairiness: very weak

Anthers - anthocyanin

Lower Glume

External surface roughness:

Extent of internal hairs: weak to medium - group 1 to 2

Beak length: short to medium

Beak shape (curvature): slightly to moderately curved

Shoulder narrow to medium

Shoulder slightly sloping or straight

Lower Lemma

Beak shape:

Grain

Colour red
Colouration with phenol: medium

Straw

Length:

Wall thickness @ 20cms below ear: very thin

Glaucosity

Flag leaf - sheath: medium to strong

Flag leaf - blade (lower side): medium

Culm neck: medium to strong
Ear - Glaucosity: medium to strong

Plant growth characters

Coleoptile-anthocyanin absent to very weak Plant - growth habit: semi-prostrate Time ear emergence: medium to late

Flag leaf - Anthocyanin colouration of auricles:



SY GRIFFIN

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs present

Awns or scurs - distribution

Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong
Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Flag leaf - anthocyanin of auricles

Coleoptile - anthocyanin colouration absent or very weak Plant - growth habit intermediate
Time of ear emergence late to very late



SY INSITOR

CHARACTERS IN DETAIL

Ear

white Colour Length medium Density lax to medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2 Awns or scurs - length long to very long Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium
Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong to very strong Flag leaf- blade (lower side) strong to very strong

Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



SY LOKI

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length short to medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence medium to late



SY MEDEA

strong

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/4Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs strong to whole surface
Beak length very short to short
Beak curvature slightly curved
Shoulder width medium to broad

Shoulder shape elevated

Lower lemma

Beak curvature

Grain

Colour red

Colouration with phenol dark to very dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath absent or very weak Flag leaf- blade (lower side) absent or very weak Culm - neck absent or very weak Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium



SY MONZA

CHARACTERS IN DETAIL

Grain

Colour reddish Colouration with phenol dark

Plant

Anthocyanin colouration of coleoptile very weak to weak

Growth habit intermed. to semi-prostrate

Frequency of plants with recurved flag leaves absent or very low Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late Length (stem, ears, awns or scurs) short Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong Flag leaf blade (lower side) medium to strong

Ear strong Culm - neck strong

Lower glume

Hairiness of external surface absent

Shoulder width very narrow to narrow Shoulder shape horizontal to slightly elevated

Length of beak medium

Shape of beak slight to moderately curved Extent of internal hairs very large (Group 3)

white

External surface roughness smooth

Ear

Density dense Length (excluding awns or scurs) medium Awns or scurs - presence scurs present Awns or scurs - length long to very long

Colour

Shape in profile parallel sided Apical rachis hairiness on convex surface small upper 1/2

Awns or scurs - distribution



THEODORE

CHARACTERS IN DETAIL

Ear

white Colour Length medium dense Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/4 Awns or scurs - length medium to long Apical rachis segment - convex face hairiness very weak to weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs weak to medium (group 1+)

Beak length medium
Beak curvature strongly curved
Shoulder width narrow
Shoulder shape elevated

Lower lemma

Beak curvature geniculate

Grain

Colour red

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



TIMARU

CHARACTERS IN DETAIL

Ear

Colour white
Length long
Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution tip only
Awns or scurs - length: very short

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong
Culm - neck medium to strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit semi-erect to intermediate

Time of ear emergence late



TOBY

CHARACTERS IN DETAIL

Ear

Colour white Length short

Density dense to very dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium to strong (group 2+)

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol medium

Straw

Length short Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong to very strong
Flag leaf- blade (lower side) strong to very strong
Culm - neck strong to very strong
Ear absent or very weak

Plant growth characters

Coleoptile - anthocyanin colouration very weak to weak
Plant - growth habit intermed. to semi-prostrate
Time of ear emergence late to very late

Flag leaf - anthocyanin of auricles absent or very weak



TORCH

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density medium to dense

Shape-side profile parallel sided

Awns or scurs - presence scurs present

Awns or scurs - distribution upper ¼

Awns or scurs - length: medium

Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak lengthmediumBeak curvatureslightly curvedShoulder widthnarrowShoulder shapeelevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear medium

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium



TORPHINS

CHARACTERS IN DETAIL

Ear

Colour white

Length short to medium

Density dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ¼

Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness medium

Anthers - anthocyanin colouration

Lower glume

External surface rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit intermediate
Time of ear emergence medium to late



TREASURE

absent

CHARACTERS IN DETAIL

Anthers - anthocyanin colouration

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthlongApical rachis segment - convex face hairinessmedium

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long
Beak curvature strongly curved
Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour red
Colouration with phenol very dark

Straw

Length very short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath very strong

Flag leaf- blade (lower side) strong to very strong

Culm - neck very strong
Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium
Plant - growth habit intermediate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



TRENDY

CHARACTERS IN DETAIL

Grain

Colour reddish
Colouration with phenol medium

Plant

Anthocyanin colouration of coleoptile very weak to weak
Growth habit intermediate
Frequency of plants with recurved flag leaves
Anthocyanin colouration of flag leaf auricles absent or very weak

Time of ear emergence late
Length (stem, ears, awns or scurs) short
Straw wall thickness (20cm below ear) thin

Glaucosity

Flag leaf sheath strong to very strong

Flag leaf blade (lower side) strong
Ear strong

Culm - neck strong to very strong

Lower glume

Hairiness of external surface absent Shoulder width narrow

Shoulder shape
Length of beak
Shape of beak
Shape of beak
Strongly curved

Extent of internal hairs very large (Group 3)

External surface roughness rough

Ear

Density medium
Length (excluding awns or scurs) medium
Awns or scurs - presence scurs present
Awns or scurs - length medium to long

Colour white
Shape in profile parallel sided
Apical rachis hairiness on convex surface small to medium
Awns or scurs - distribution upper 1/4



TUXEDO

CHARACTERS IN DETAIL

Ear

Colour white Length medium

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ¼

Awns or scurs - length: medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak lengthmediumBeak curvatureslightly curvedShoulder widthnarrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong

Culm - neck medium to strong Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

Time of ear emergence medium to late



TWISTER

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ½

Awns or scurs - length: long to very long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2
Beak length medium to long

Beak curvature moderate to strongly curved

Shoulder width narrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium
Culm - neck medium
Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium



UNIVERSAL

CHARACTERS IN DETAIL

Ear

white Colour

medium to long Length Density lax to medium Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper 1/2

Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth Extent of internal hairs medium (2) Beak length medium to long Beak curvature strongly curved

Shoulder width medium

Shoulder shape straight to elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red Colouration with phenol dark

Straw

short to medium Length Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong Flag leaf- blade (lower side) medium to strong Culm - neck medium to strong Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence late



VELOCITY

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributiontip only

Awns or scurs - length short to medium

Apical rachis segment - convex face hairiness medium
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)
Beak length medium to long

Beak curvature strongly curved to geniculate

Shoulder width medium to broad

Shoulder shape strongly elevated with 2nd pt. present

Lower lemma

Beak curvature geniculate

Grain

Colour red
Colouration with phenol dark

Straw

Length short to medium Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) weak to medium

Culm - neck medium Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence early to medium



VERSO

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density medium
Shape-side profile tapering
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 3/4
Awns or scurs - length long

Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium (group 2)

Beak length medium

Beak curvature v.slightly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature straight to slightly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side) medium to strong

Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration weak
Plant - growth habit intermediate
Time of ear emergence medium to late



VISCOUNT

CHARACTERS IN DETAIL

Ear

Colour white
Length medium
Density dense
Shape-side profile parallel s

Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ¼
Awns or scurs - length: medium to long

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak length medium

Beak curvature moderately curved

Shoulder width medium
Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) medium

Culm - neck medium to strong

Ear medium

Plant growth characters

Coleoptile - anthocyanin colouration strong

Plant - growth habit semi-erect to intermediate

late

Time of ear emergence



Variety Name: Vulkan

crop: Winter Wheat

Present

Breeder's Reference: AFP No:

Origin: Date: 11 March 2014

Code: VULK

Coleoptile-anthocyanin colouration:

*Plant - growth habit: Semi erect to intermediate

*Time ear emergence: Very early to early

Flag leaf - Anthocyanin colouration of auricles:

Frequency of plants with recurved flag leaves: Low
*Flag leaf - Glaucosity of sheath: Medium

Flag leaf - Glaucosity of leaf blade: Very weak to weak

Anthers - anthocyanin colouration:

Glaucosity of neck: Weak to medium

Hairiness of uppermost node:

*Plant - height: Medium

*Straw - wall thickness @ 20cms below ear: Very thick or filled

*Ear - Glaucosity:

Ear - Supernumerary spikelets - frequency:

*Ear - Colour: White

*Ear - Shape (side profile): Parallel sided

Ear - Length: Very short

*Ear - Density: Medium

*Awns or scurs - presence: Awns or scurs - distibution:

*Scurs at tip of ear - length: Medium
*Awns at tip of ear - length: Medium

Lower glume - Beak length:

Long to very long

Lower glume - Beak shape (curvature):

Slightly curved

Lower glume - Beak tip:

Lower glume - Shoulder width: Absent or very narrow

Lower glume - Shoulder type: Sloping

Lower glume - External surface roughness:

Lower glume - Extent of internal hairs: Medium

Lower glume - Internal imprint: Lower lemma - Beak shape: Lower lemma - Beak length-: Lower lemma - Beak swelling:

Apical rachis segement - convex face hairiness:

*Grain - colour:

Red

Grain - Shape:

Grain - Brush hair length in dorsal view:

Grain - Colouration with phenol:

*Seasonal type: Winter

Extra characters:



WARRIOR

CHARACTERS IN DETAIL

Ear

Colour white Length short

Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper 1/4
Awns or scurs - length medium to long
Apical rachis segment - convex face hairiness weak to medium

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough

Extent of internal hairs weak to medium - Group 1+

Beak lengthmediumBeak curvatureslightly curvedShoulder widthnarrow to medium

Shoulder shape elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear n/a

Glaucosity

Flag leaf - sheath strong
Flag leaf - blade (lower side) strong
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration medium to strong
Plant - growth habit semi-erect
Time of ear emergence medium to late



WEAVER

strong

CHARACTERS IN DETAIL

Ear

Colour white

Length medium to long
Density medium to dense
Shape-side profile parallel sided
Awns or scurs - presence scurs present
Awns or scurs - distribution upper ¼
Awns or scurs - length: long

Apical rachis segment - convex face hairiness

Anthers - anthocyanin colouration

Lower glume

External surface slightly rough
Extent of internal hairs medium - Group 2

Beak length medium
Beak curvature strongly curved

Shoulder width narrow Shoulder shape elevated

Lower lemma

Beak curvature strongly curved

Grain

Colour red
Colouration with phenol very dark

Straw

Length short to medium

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong
Flag leaf- blade (lower side) strong
Culm - neck strong

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration absent to weak Plant - growth habit intermediate Time of ear emergence medium



WELCOME

CHARACTERS IN DETAIL

Ear

Colour white

Lengthshort to mediumDensitymedium to denseShape-side profileparallel sidedAwns or scurs - presenceawns presentAwns or scurs - distributionwhole length

Awns or scurs - length very long
Apical rachis segment - convex face hairiness absent or very weak

Anthers - anthocyanin colouration absent

Lower glume

External surface smooth
Extent of internal hairs medium (2)

Beak length very long
Beak curvature moderately curved
Shoulder width narrow

Shoulder shape elev. to elev with 2nd point

Lower lemma

Beak curvature straight

Grain

Colour red
Colouration with phenol very dark

Straw

Length medium

Wall thickness @20cms below ear absent or very thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side) medium
Culm - neck medium

Ear medium to strong

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium to late



WELFORD

CHARACTERS IN DETAIL

Ear

ColourwhiteLengthmediumDensitymediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2Awns or scurs - lengthmedium

Apical rachis segment - convex face hairiness medium to strong

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature moderately curved

Shoulder width narrow

Shoulder shape straight to elevated

Lower lemma

Beak curvature moderately curved

Grain

Colour red
Colouration with phenol dark

Straw

Length short Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong

Ear weak to medium

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermed. to semi-prostrate

Time of ear emergence medium

Flag leaf - anthocyanin of auricles absent or very weak



XI19

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs present

Awns or scurs - length:shortAwns or scurs - distributiontip onlyApical rachis segment - convex face hairinessmediumAnthers - anthocyanin colourationabsent

Lower glume

External surface smooth

Extent of internal hairs strong - Group 3
Beak length medium

Beak curvature moderately curved

Shoulder width narrow Shoulder shape straight

Lower lemma

Beak curvature moderately to strongly curved

Grain

Colour red
Colouration with phenol dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong

Flag leaf- blade (lower side)

Culm - neck strong

Ear strong to very strong

Plant growth characters

Coleoptile - anthocyanin colouration absent or very weak
Plant - growth habit intermed. to semi-prostrate

Time of ear emergence late
Flag leaf - anthocyanin of auricles very weak



ZEBEDEE

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitylax to mediumShape-side profileparallel sidedAwns or scurs - presencescurs presentAwns or scurs - distributionupper ¼

Awns or scurs - length: short to medium

Apical rachis segment - convex face hairiness weak
Anthers - anthocyanin colouration absent

Lower glume

External surface smooth

Extent of internal hairs medium - Group 2

Beak length medium

Beak curvature slight to moderately curved

Shoulder width narrow to medium

Shoulder shape straight

Lower lemma

Beak curvature slightly to moderately curved

Grain

Colour red

Colouration with phenol light to medium

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium to strong

Flag leaf- blade (lower side)

Culm - neck medium to strong

Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration medium

Plant - growth habit intermediate

Time of ear emergence medium to late

Flag leaf - anthocyanin of auricles absent or very weak



ZOOM

CHARACTERS IN DETAIL

Ear

Colour white

Lengthmedium to longDensitymediumShape-side profiletaperingAwns or scurs - presencescurs presentAwns or scurs - distributionupper 1/2

Awns or scurs - length long to very long
Apical rachis segment - convex face hairiness medium to large

Anthers - anthocyanin colouration absent

Lower glume

External surface slightly rough Extent of internal hairs strong (group 3)

Beak curvature strongly curved

Shoulder width narrow

Shoulder shape slightly elevated to strongly elevated

Lower lemma

Beak curvature strongly curved to geniculate

Grain

Colour reddish

Colouration with phenol medium to dark

Straw

Length very short to short

Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath strong to very strong

Flag leaf- blade (lower side) strong
Culm - neck strong
Ear strong

Plant growth characters

Coleoptile - anthocyanin colouration weak

Plant - growth habit semi-prostrate

Time of ear emergence late

Flag leaf - anthocyanin of auricles absent or very weak



ZULU

CHARACTERS IN DETAIL

Ear

white Colour Length medium medium Density Shape-side profile parallel sided Awns or scurs - presence scurs present Awns or scurs - distribution upper ¾ Awns or scurs - length: long Apical rachis segment - convex face hairiness medium absent Anthers - anthocyanin colouration

Lower glume

External surface smooth
Extent of internal hairs strong - Group 3

Beak lengthmediumBeak curvatureslightly curvedShoulder widthnarrowShoulder shapeelevated

Lower lemma

Beak curvature slightly curved

Grain

Colour red Colouration with phenol dark

Straw

Length medium Wall thickness @20cms below ear thin

Glaucosity

Flag leaf - sheath medium
Flag leaf- blade (lower side) weak to medium
Culm - neck medium

Ear

Plant growth characters

Coleoptile - anthocyanin colouration weak to medium
Plant - growth habit intermed. to semi-prostrate

medium

Time of ear emergence medium to late

